

**Lower Cretaceous reservoir development in the North Sea Central Graben and potential analogue settings in the Southern Permian Basin and South Viking Graben**

**Frank Zwaan<sup>1,2\*</sup>**

<sup>1</sup> Project work: VU University Amsterdam, Faculty of Earth and Life Sciences, De Boelelaan 1085-1087, 1081 HV Amsterdam, <sup>2</sup> Currently: University of Bern, Institute of Geological Sciences, Baltzerstrasse 1+3, CH-3012 Bern, Switzerland)

\*Corresponding author: [frank.zwaan@geo.unibe.ch](mailto:frank.zwaan@geo.unibe.ch)

**ABSTRACT**

Much of the future hydrocarbon exploration potential in the North Sea lies in locating stratigraphic traps and discrete reservoir intervals. This study assesses the potential for Lower Cretaceous reservoirs, with particular focus on the Norwegian Central Graben and proposed methods to identify future prospects over a wider area. Seismic interpretation and well data reveal the structure and sedimentology of the study area. Although the region was isolated from a large hinterland in the Early Cretaceous, potential local sediment sources, sediment transport routes and areas with possible reservoir development are identified. The greater Mandal High area, where Lower Cretaceous shoreface deposits and submarine fan systems are postulated, is suggested for primary focus. Similar deposits may have developed around the other exposed highs in the region, although several were drowned towards the end of the Early Cretaceous. Detailed seismic and stratigraphic analysis will be necessary to identify individual reservoir units. Since similar settings may have occurred in the adjacent South Viking Graben and Southern Permian Basin regions during the Early Cretaceous, further reservoir assessment is recommended for the North Sea in general.

**KEYWORDS:** Lower Cretaceous, Tectonics, Reservoir prediction, Hydrocarbon exploration

Abstract Word Count: 180

Main Body Word Count (excluding abstract, figure captions & references): 7723

References Word Count: 3999, including DOIs

Figure Caption Word Count: 1400 Words (incl table caption), 11 figures, 1 table

**THIS IS A POSTPRINT VERSION, FINAL VERSION PUBLISHED IN:**  
KILHAMS, B., KUKLA, P. A., MAZUR, S., MCKIE, T., MIJNLIEFF, H. F. & VAN OJIK, K. (eds) Mesozoic Resource Potential in the Southern Permian Basin. Geological Society, London, Special Publications, 469, <https://doi.org/10.1144/SP469.3> © 2018

The Central Graben of the North Sea represents a prolific and mature hydrocarbon province (e.g. Brooks 1990). Despite 50 years of exploration, further potential continues to be unlocked as demonstrated by the Upper Jurassic shoreface play (e.g. Edvard Grieg and Johan Sverdrup discoveries at the Norwegian Utsira High, Jørstad 2012; NPD 2017; Lundin 2017) and by the Upper Jurassic turbidite play in the United Kingdom (UK) Moray Firth (e.g. Buzzard field, Fraser *et al.* 2003; Ray *et al.* 2010). This paper describes further potential in another stratigraphic horizon, namely the Lower Cretaceous reservoirs of the North Sea. Here shoreface and deep water reservoir units are developed associated with discrete Mesozoic rifting phases and related localised depocenters.

Lower Cretaceous deep marine sandstones represent an important play in the UK Moray Firth (e.g. Garrett *et al.* 2000; Johnson *et al.* 2005, Figs. 1a, 2), where large amounts of sand from the exposed East Orkney High and Halibut Horst were shed into adjacent basins of the Inner- and Outer Moray Firth, forming the reservoirs for various hydrocarbon fields (e.g. Scapa, Britannia, McGann *et al.* 1991; Ainsworth *et al.* 2000). Various studies have established sediment transport directions (Hailwood & Ding 2000), sediment provenance areas (Blackbourn & Thomson 2000) and a sequence stratigraphic framework (Jeremiah 2000) in this area.

In contrast to the Moray Firth, the Lower Cretaceous play remains underdeveloped in the Central Graben (Copestake *et al.* 2003; NPD 2017). While the generalised Mesozoic sequence is visible on seismic in the Moray Firth, the thick (2000 to 4500 m) Cenozoic and Upper Cretaceous overburden in the Central Graben presents an issue for seismically driven reservoir identification within the Lower Cretaceous (up to 30 m net reservoir thickness) (Argent *et al.* 2000; Law *et al.* 2000). Further factors impeding seismic assessment include multiples (induced by the base of the overlying Upper Cretaceous chalk units) and the general low impedance contrast between sandstones and shales in this interval (Oakman 2005). Although seismic imaging quality has improved considerably in recent years (e.g. Hampson *et al.* 2010), alternative methods are required to assess the Lower Cretaceous units and to locate potential prospects. Attempts to delineate reservoir development have been made in the UK Central Graben (UKCG) with the use of regional 3D seismic and well data (Milton-Worsell *et al.* 2006, Fig. 1a), indicating significant potential for Lower Cretaceous sandstone development. Currently, other than local studies (e.g. Rossland *et al.* 2013), no published work has established a similar overview of Lower Cretaceous reservoir potential in the Norwegian sector of the Central Graben.

This study, therefore, aims to assess the potential for Lower Cretaceous sand bodies in the Norwegian Central Graben (NCG) and to link this interpretation to adjacent areas in the UK and to the Danish, German and Dutch parts of the North Sea Rift System (Southern Permian Basin area), as well as to the South Viking Graben (Fig. 1a).

## GEOLOGICAL SETTING

The geological history of the Central North Sea has generated a diverse stratigraphic record, which is hereby described utilising a Norwegian stratigraphic nomenclature (Isaksen & Tonstad 1989; NPD 2017, Fig. 2). The oldest known units in the Norwegian Central Graben are of pre-Permian age; Silurian to Devonian metamorphic basement overlain by Old Red sandstones (Fossen *et al.* 2008). These units are followed, in stratigraphic order, by varied Carboniferous deposits (NPD 2014), Lower Permian Rotliegend sandstones and thick Upper Permian Zechstein evaporites plus dolomites. The halite sequences of the Zechstein have had a profound influence on the tectonic style; decoupling underlying and overlying strata (Hodgson *et al.* 1992; Stewart 2007; Ge *et al.* 2016; Jackson & Lewis 2016; Van Winden this volume). Lower Triassic Smith Bank Shales and Middle to Upper Triassic Skagerrak Sandstone deposition coincided with Late Triassic faulting along the inherited Caledonian structural grain (Bartholomew *et al.* 1993; UKDD 2007). This first rifting phase formed the general Central Graben structure, as these Triassic faults were partially reactivated in the Late Jurassic and Early Cretaceous (Rathey & Hayward 1993), although the main Triassic and Jurassic to Early Cretaceous depocenters do not precisely coincide (Erratt *et al.* 1999). Uplift and erosion due to an Early Jurassic mantle plume formed the Mid-Cimmerian unconformity in the Central North Sea (Underhill & Partington 1993). Middle Jurassic Bryne Formation coastal plain deposits succeed the hiatus (Bergan *et al.* 1989), followed by a second erosional surface. Eventual dome collapse coincided with the onset of renewed extension in the Northern North Sea (Graversen 2006), propagating into the Central North Sea during Late Jurassic times (Rathey & Hayward 1993). As rifting proceeded, sediment-starved deep marine basins developed (Copestake *et al.* 2003). This transgression is recorded in the syn-rift Jurassic Tyne Group by the Ula Formation shoreface sandstones and subsequent deep marine shales, including Mandal Formation source rock (Gautier 2005; Nøttvedt & Johannessen 2008).

Major extension ceased towards the end of the Kimmeridgian (Milton 1993), although a secondary phase of rift activity may have continued into the Earliest Cretaceous or Ryazanian in the Norwegian area, especially in the NCG (Gowers *et al.* 1993; Sears *et al.* 1993; Zanella *et al.* 2003; Ge *et al.* 2016). Rifting ceased as extensional stresses shifted to the proto-North Atlantic (Coward *et al.* 2003, Oakman 2005). Subsequently, post-rift thermal sag initiated and sediments began to cover the rift topography above the Base Cretaceous Unconformity (BCU) (Rathey & Hayward 1993). The first of these, the Cromer Knoll Group, contains mostly shales but also marly limestones (Tuxen Formation) plus shoreface to deep marine sandstones (Ran Sandstone units, Isaksen & Tonstad 1989). The sandstones represent potential reservoir bodies but their spatial and temporal extent is poorly constrained, being encountered in only a few wells. In the Aptian, another shift in tectonics and oceanography, the “Austrian event”, occurred. This coincided with the onset of alpine compression and the opening of the North Atlantic, leading to more restricted basins and to the deposition of dark muds of the Sola Formation, overlain by calcareous Rødby Formation sediments (Garrett *et al.* 2000; Copestake *et al.* 2003).

Global sea level rise and a shift to a tropical climate in the Late Cretaceous saw the development of massive Upper Cretaceous chalk units in the North Sea sag basin (Surlyk *et al.* 2003). Thermal subsidence was, however, interrupted by local inversion pulses and associated Zechstein salt diapirism (Cartwright 1989; Johnson *et al.* 2005; Van Winden this volume). Renewed sediment input from the eroding North Atlantic rift shoulders gave rise to widespread turbidite systems in the Paleocene and Eocene. From the Oligocene onwards, thermal sag continued, concentrated in the NW of the NCG (Gowers & Sæbøe 1985), while the North Sea basin gradually filled in with thick clastic sequences.

## ESTABLISHED LOWER CRETACEOUS UK RESERVOIRS

Three major sequences of sandstone deposits occur in the Inner Moray Firth (Copestake *et al.* 2003, Fig. 2); Ryazanian-Valanginian Punt sandstones SW of the Halibut Horst, Wick Sandstones south of the East Orkney High and Scapa Sandstones east of the Halibut High (locations in Fig. 1). In Barremian times, Coracle Sandstones of the Wick Fm occurred south of the East Orkney High and the Halibut High, whilst Scapa Sandstones were still present in the Witch Ground Graben (Jeremiah 2000). These units of the Lowermost Cretaceous (Ryazanian-Barremian play) were deposited during a phase of low sea level due to tectonic activity related to Austrian compression, ending with a major flooding event in the Barremian (Crittenden *et al.* 1997; Oakman 2005). Rejuvenated tectonic activity associated with the opening of the North Atlantic led to renewed sediment influx in the Aptian (Oakman 2005). During this phase, the Kopervik fairway was established (Law *et al.* 2000) along which large amounts of sand were transported from the East Orkney High to the outer Moray Firth, where the Britannia Field is situated (Ainsworth *et al.* 2000), before an Albian transgression diminished sand influx (Oakman & Partington 1998; Jeremiah 2000).

These deposits comprise of deep marine sandstones exhibiting a variety of depositional styles including hanging-wall slope-apron fans, linear channel complexes as part of a minibasin spilling system, or localised mass flow deposits and mud-dominated slurry-flow deposits (Jones *et al.* 1999; Argent *et al.* 2000). These sedimentary systems demonstrate a high degree of complexity regarding source and transport mechanisms (Eggenhuisen *et al.* 2010). Deposition was strongly influenced by the two Early Cretaceous tectonic phases mentioned above which uplifted and exposed highs and fault scarps, as documented to the north and northwest of the main depocenters (e.g. Halibut Horst, East Orkney High, O'Driscoll *et al.* 1990; Copestake *et al.* 2003; Jeremiah 2000, Fig. 1a). Tectonic activity furthermore modified the region's bathymetry and redirected sediment transport fairways (Jeremiah 2000; Aas *et al.* 2010). These deep marine sandstones represent Lower Cretaceous reservoirs in stratigraphic or combination structural/stratigraphic traps in for instance the Britannia, Scapa and Captain fields (McGann *et al.* 1991; Jones *et al.* 1999; Pinnock *et al.* 2003).

Although the Forties-Montrose High and Marnock Terrace formed barriers that separated the UKCG depocenters from the Moray Firth during the Early Cretaceous (Fig. 1a), it is possible to extend the Lower Cretaceous Moray Firth reservoir intervals into the UKCG (Milton-Worssell *et al.* 2006) where various wells encounter Lower Cretaceous sands. This well data, in combination with seismically-derived maps, allowed Milton-Worssell *et al.* (2006) to

postulate the distribution of (mainly deep marine sandstone) bodies, sourced by the Western Platform, Forties-Montrose High and Jæren High, for both a Latest Ryazanian-Barremian play and the Aptian-Albian play in the UKCG. These plays are separated by the Fischschiefer Bed (Fig. 2), an organic-rich mudstone deposited during the Barremian flooding event, that is a regional seismic marker (Ainsworth *et al.* 2000). In this study, a similar division has been made between a “Latest Ryazanian” interval (near BCU-level) and an “Aptian-Albian” interval (near Top Lower Cretaceous level) to extend the scope into the NCG and to acquire a North Sea-wide overview (Fig. 2).

## DATA AND METHODS

2D and 3D seismic datasets, provided by Shell Upstream International, combined with data from 474 wells were used to establish a structural framework in the study area (Fig. 1). The 3D data (extent: 11,400 km<sup>2</sup>) are a compilation of the Norwegian Carmot dataset, covering the NCG, and part of the UK Megamerge dataset, covering a limited area part of the southern UKCG (Fig. 1). The quality is variable but typically consists of dominant frequencies of ca. 20-30 Hz, a wavelength of ca. 60 ms two-way travel time (TWT) with a resulting seismic resolution of ca. 15 ms TWT. This corresponds to ca. 30 m vertical resolution assuming an interval velocity of 3500 m/s. Water depths range from 40 to 100 m.

Data concerning 319 Norwegian wells in the study area were obtained from the Norwegian Petroleum Directorate (NPD) Factpages (NPD 2017). Additional well data for the UK and Danish sectors (119 and 41 wells, respectively) are from released well log and completion reports, well logs in the Shell archive (e.g. Boirie & Jeannou 1984; Statoil 1991) and published material (e.g. Isaksen & Tonstad 1989, see table 1). Additional occurrences of Lower Cretaceous sandstones in UKCG wells are adopted from Milton-Worrrell *et al.* 2006).

The following regional seismic horizons were mapped in two way time (TWT) on 3D seismic and calibrated with time-converted (via well checkshots and calibrated sonic logs) lithostratigraphically defined well tops from the Shell database and the NPD (Fig. 2):

- Base Cenozoic, (64 Ma);
- Top Lower Cretaceous (100 Ma);
- Base Cretaceous Unconformity (BCU, 140 Ma);
- Top Rotliegend (270 Ma).

Milton-Worrrell *et al.* (2006) mapped the Fischerbank Schiefer Bed, which defines the boundary between their two Lower Cretaceous plays (Fig. 2). In this study this marker could not be traced due to a lack of accurately constrained well picks. The time maps of the four interpreted seismic horizons are combined with existing digital TWT seismic horizon maps provided by Shell Upstream International that allow an extension of the survey further into British and Danish territorial waters (study area, Fig. 1). A time difference assessment between the seismic horizons yields isochron maps, illustrating where the thickest sequences within the Cenozoic, Upper Cretaceous and Lower Cretaceous intervals are situated, revealing the general structural trends in the study area (Figs. 3 and 4). Due to the large study

area, no time to depth conversion was carried out which means these structural trends are somewhat qualitative.

Two more lithostratigraphically-constrained horizons have been mapped in TWT on five additional regional 2D seismic transects to provide an additional link to previous studies (S1-S5, Figs. 1, 2, 5):

- Base Upper Jurassic (ca. 165 Ma);
- Top Zechstein (252 Ma).

Although the available seismic coverage does not include Denmark, an earlier study (Møller & Rasmussen 2003) provides a useful additional transect across the Danish border (S6, Fig. 5f). In combination with the seismic horizon time and isochron maps, these transects offer a detailed insight into the structural framework of the extended study area, revealing the locations of the main basins, highs, diapirs and faults (Figs. 3-5). Subsequently, the results of the seismic interpretation are integrated with published data from Copestake *et al.* (2003), Japsen *et al.* (2003), Milton-Worsell *et al.* (2006) and Rossland *et al.* (2013) for an assessment of Lower Cretaceous reservoir potential in the extended study area, of which well data provide a first impression (Figs. 4 and 6).

The basic methodology applied by Milton-Worsell *et al.* (2006) has been adopted. Combined isochron maps of the extended study area indicate zones with thin Lower Cretaceous deposits, which were potentially exposed and prone to erosion during the Early Cretaceous (Fig. 7). At these places, well data provides the true thickness of the Lower Cretaceous sequence and the lithology in subcrop below the BCU. Devonian metamorphic rocks and volcanics, Rotliegend, Triassic Skagerrak, Middle Jurassic Bryne and Upper Jurassic Ula sandstones (Fig. 2) in subcrop indicate whether a specific locality was part of a potential sand source area during the Earliest Cretaceous. The presence of sand provenance areas is considered the most important factor controlling sandstone development since the Early Cretaceous was dominated by pelagic mud deposition (Fig. 2). This exercise is repeated for the Aptian-Albian reservoir interval, where the sand-prone lithologies in subcrop below the Top Lower Cretaceous horizon are charted (Fig. 8). The isochron maps subsequently allow the tracing of possible sediment transport fairways, by interpreting depocenters as drainage areas and barriers separating them as watersheds. Sediment transport is assumed to have followed the bathymetry given by the isochron maps, leading sediments from the highs to the depocenters. Thus, combining the isochron map, drainage and sand source areas; potential sand transport routes for the Latest Ryazanian and the Aptian-Albian are mapped (Figs 7 and 8). Well data allows a qualitative check of these interpretations: where sandstones occur in wells, a plausible link with a nearby sand source area can be inferred. If no such well data is available, sediment transport between source and depocenter remains speculative. It is recognised that the sandstones recorded in these wells are not necessarily linked to the postulated source areas and that those links would need to be proven via further investigation involving advanced seismic and well analysis techniques that are beyond the scope of this study.

235

**STRUCTURAL FRAMEWORK INTERPRETATION**

236 In general, a series of NNW-SSE orientated en-echelon (rift) basins, normal faults and tilted  
 237 fault blocks follow the larger NW-SE Central Graben trend (Fig. 1, 3-5). The NCG structure  
 238 is bounded by the Sørvestlandet High and the Ringkøbing-Fyn High to the east and by the  
 239 Mid North Sea High to the south-west (Figs. 3-4).

240 The main depocenters, as identified on the isochron maps, are situated in the Breiflab Basin  
 241 in the NW (Fig. 4a, 5, S1), where up to 8 km of subsidence has occurred (NDD 2012).  
 242 However, the locations of these depocenters do not coincide with the thickest Upper Jurassic  
 243 deposits in the SE part of the Feda Graben, Søgne Basin and Gertrud Graben (Erratt *et al.*  
 244 1999, Fig. 5, S5, S6). This discrepancy is a result of later differential thermal subsidence and  
 245 sediment infill (Gowers & Sæbøe 1985). Normal faults are omnipresent in the area, but major  
 246 differences in structural style occur between the Pre-Zechstein units, Triassic, Upper Jurassic,  
 247 Lower Cretaceous syn-rift strata and post-rift infill. The Josephine High (Fig. 5, S1), Hidra  
 248 High (Fig. 5, S2), Border High (Fig. 5, S4), Mandal High (Fig. 5, S5, S6) Cod Terrace (Fig.  
 249 5, S1) and Piggvar Terrace (Fig. 5, S5) represent Pre-Zechstein basement blocks forming  
 250 major structural highs or terraces. Several large salt domes occur within the area (e.g. Fig. 5,  
 251 S1).

**252 Late Jurassic-Early Cretaceous rift structures**

253 Due to Mid-Jurassic thermal doming and associated erosion, few Lower Jurassic units are  
 254 preserved in the study area. In contrast, significant Upper Jurassic sediments, recording the  
 255 latest North Sea rift phase, occur locally in extensional basins. These units are best developed  
 256 in the south of the study area, where the Feda Graben, Gertrud Graben and Søgne Basin half-  
 257 graben accommodate some 2 km of Upper Jurassic sequences (Fig. 5, S5, S6) as part of the  
 258 large-scale left-stepping en-echelon Central Graben structure (Erratt *et al.* 1999, Fig. 1a).  
 259 Many Triassic faults affect Upper Jurassic strata, indicating fault reactivation, e.g. the  
 260 Skrubbe Fault and Coffee Soil Fault bounding the Feda Graben and Søgne Basin,  
 261 respectively (Fig. 5, S5, S6). Rifting caused salt movement and diapirism which impacted  
 262 Upper Jurassic sedimentation e.g. in the Søgne Basin.

263 Subsequently, the major Lower Cretaceous deposits are shifted westward compared to the  
 264 Upper Jurassic depocenters (Figs. 5, S1, S3, S4). A distinct feature is the Early Cretaceous  
 265 reactivation of the Pre-Zechstein half-graben west of the Border High, where Upper Jurassic  
 266 or Triassic units are absent (Fig. 5, S4). Also striking is the lack of Early Cretaceous tectonic  
 267 activity in the Søgne Basin; in contrast to significant Triassic and Upper Jurassic syn-tectonic  
 268 units, little to no Lower Cretaceous sediments occur (Fig. 4b, 5, S5, S6).

269 The character of the Early Cretaceous basins varies considerably. The Border High and  
 270 Breiflab Basins are fault-bounded and show thickening towards the boundary faults,  
 271 indicating syn-rift deposition (Fig. 5, S1, S4). Other rift-bounded basins are found west of the  
 272 Hidra High (Fig. 5, S2), at well NO 2/4-10 (Fig. 5, S3) and west of the Mandal High (Fig. 5,  
 273 S6). Yet the filling of pre-existing deep underfilled Jurassic basins as well as sediment  
 274 compaction effects could partially account for these observations (Rathey & Hayward 1993;

Coward *et al.* 2003). At various localities, salt motion affected Early Cretaceous deposition: e.g. above the Hydra High (Fig. 5, S2) and at well NO 2/4-3 (Fig. 5, S4). In other parts of the study area, depocenters exhibit sag-type geometries, e.g. east of well UK 30/17B-3 and above the Hydra High (both in Fig. 5, S2), west of the NO 1/6-5 diapir (Fig. 5, S3) and in the Ål Basin (Fig. 5, S5, S6). Faults do not generally continue to the top of the Early Cretaceous, except for those associated with later tectonic inversion.

As such, cessation of rifting is shown to be diachronous. The westward shift of the Early Cretaceous depocenters with respect to the Jurassic rifts might indicate a change in extensional regime near the start of the Cretaceous, as proposed by previous authors (e.g. Erratt *et al.* 1999), before extension activity ceased altogether due to the opening of the young North Atlantic (Rattee & Hayward 1993).

### **Post-rift and tectonic inversion structures**

The Late Cretaceous and Cenozoic units dominantly show gentle sag geometries along the NW-SE trend of the NCG, indicating further post-rift thermal subsidence. At the Breiflabb basin on the UK/Norwegian border, thermal subsidence was strongest creating a major Late Cretaceous/Cenozoic depocenter (Figs. 4a, 5, S1, 6a, Gowers & Sæbøe 1985). However, signs of inversion are also noted, for instance at the Lindesnes Ridge where Early Cretaceous syn-rift deposits are uplifted along Skrubbe Fault, (Figs. 5, S1, 6e). Inversion-related structures (inverted grabens and diapirs/salt domes) disturb not only the Upper Cretaceous deposits, but also Cenozoic strata (Figs. 3-5), indicating multiple inversion phases (Gowers *et al.* 1993).

## **LOWER CRETACEOUS RESERVOIR INTERPRETATION**

### **Sandstone occurrences in Norwegian and Danish wells**

In contrast to the UKCG, where numerous wells encounter Lower Cretaceous sandstones (Milton-Worssell *et al.* 2006), only three wells in the NCG area (from a total of 160 Lower Cretaceous penetrations) are reported to contain similar deposits (NPD 2017, Figs. 4b, 6). The sandstones in these wells are lithostratigraphically defined as Ran Sandstone units (NPD 2017) and, in contrast with the deep marine character of most equivalent Lower Cretaceous sandstones in the UK, are interpreted as shallow submarine fans (Isaksen & Tonstad 1989; Milton-Worssell *et al.* 2006).

Well NO 2/1-8 on the Cod Terrace contains a 4 m interval of Ran Sandstones, but no further details on lithology, or reservoir properties are publicly available (Fjellanger 1986; NPD 2013). These sandstones appear below the Hauterivian-Barremian Tuxen Fm and are, therefore, assigned to the Ryazanian reservoir interval (Fig. 2). Reference well NO 2/7-15 in the Feda Graben (Isaksen & Tonstad 1989, Fig. 4b) contains a 48 m thick Ran Sandstone sequence. Cores taken from the lowermost part of this succession are described as dominantly clay-rich siltstones with occasional micro-porosity and fractures with minor hydrocarbon shows (Phillips 1981). However, drill stem tests demonstrated the section to be tight (NPD 2017). The age of these Ran Sandstones is poorly constrained, but they are



attributed to the Albian-Aptian reservoir interval due to their occurrence directly below the Aptian-Albian Sola Fm (Isaksen & Tonstad 1989, Fig. 2).

In well NO 3/7-3, east of the Mandal High (Fig. 4b), a 107 m thick Ran Sandstone sequence occurs on top of the BCU (NPD 2017). These deposits consist of a lower unit of dolomitic and glauconitic sandstones, interbedded with dolomitic and shaley layers, and an upper unit of massive coarse-grained sandstones with occurrences of chalky, sandy limestone, capped by carbonates containing some lignite (Verolles 1982). The massive sandstones (60-70% quartz) are cemented but represent good reservoir potential with porosities and permeabilities between 20-28 % and 0.5 to 10 D respectively (Verolles 1982; Boirie & Jeannou 1984). The NO 3/7-3 Ran Sandstones were deposited as lenticular sheets or slope apron bodies in a restricted and proximal, relatively shallow marine environment (100-200 m water depth, Verolles 1982), which evolved into an open marine setting towards the end of the Early Cretaceous (Boirie & Jeannou 1984). Since the Ran Sandstones are of Ryazanian age (Boirie & Jeannou 1984), they belong to the Latest Ryazanian reservoir interval.

Four other Norwegian wells encountering Ran Sandstone are situated to the NE, in block 17, at a considerable distance from the North Sea rift basins and outside the extended study area. The implications of these sandstone occurrences will be addressed in the South Viking Graben regional overview below.

In the Tail End Graben (Denmark), 9 m thick Lower Cretaceous subangular to subrounded and poor to moderately sorted, fine grained “Kira Sandstones” are found above BCU-level in the Amalie-1 well, probably deposited as part of a submarine fan system (Statoil 1991, Fig. 6). These sandstones are oil-bearing and of excellent reservoir quality with high porosities and permeabilities (0.213 and 319 mD, respectively) and a net-to gross ratio of 0.339 (Statoil 1991). Further Latest Ryazanian sandstones, although thinner, occur in the Tabita-1, Svane-1 and Iris-1 wells south of the Amalie-1 well (Figs. 4b). The Tabita-1 “Kira Sandstone equivalent” at the base of the Lower Cretaceous contains mostly claystone with very fine grained silt- and (quartz) sandstone striae (1-3 cm), as well as cross bedding with erosional surfaces (Bonde *et al.* 1994). A core from this interval contains conglomeratic intervals of unweathered, angular clasts of metamorphic basement material, as well as folded and disturbed mudstone beds. Both facies are indicative of slope process, whilst the lack of wave-related structures in the core suggests a depositional environment below storm wave base (Bonde *et al.* 1994). In the Svane-1 well, very fine to fine grained, subrounded, poorly sorted calcareous quartz sandstones with an argillaceous matrix and net-to-gross ratios up to 0.85 are found above the BCU (Thorsrud *et al.* 2002). The Iris-1 well contains various levels of thin sandstone in the Valhall Fm overlying the BCU which are “a few” meters thick (Britoil 1985). The cored material from this well is predominantly fine-grained and similar to that in the Tabita-1 well (Bonde *et al.* 1994). Further to the west, Lower Cretaceous (Latest Ryazanian-Early Hauterivian) fine to medium grained, poorly sorted sandstones, belonging to the Latest Ryazanian reservoir interval, are present in the Sten-1 well (Kern *et al.* 1983), making a total of 5 wells encountering Lower Cretaceous sandstones in the Danish part of the study area (Fig. 4b).

## 356 Latest Ryazanian reservoir distribution

357 An interpretation of the Latest Ryazanian reservoir interval is presented in Fig. 7 and depicts  
 358 the sandstone occurrences in wells, potential source areas with sand-prone lithologies  
 359 subcropping the BCU and sediment transport fairways to depocenters identified on the Lower  
 360 Cretaceous isochron map.

361 Milton-Worrrell *et al.* (2006) demonstrated the potential for marine sandstone development  
 362 in the Ryazanian-Barremian interval of the UKCG, with the Forties-Montrose High and  
 363 Western Platform interpreted as provenance areas. Closer to the Norwegian-British border,  
 364 sand-prone lithologies are found in subcrop below the BCU at the Josephine High (Skagerrak  
 365 Fm), Auk Ridge (Rotliegend) and Argyll Field at the Mid North Sea High (Rotliegend, Ula  
 366 Fm and Skagerrak Fm). These represent potential sand source areas for the surrounding  
 367 depocenters where multiple well penetrations occur (Milton-Worrrell *et al.* 2006). The Auk  
 368 Ridge is also the likely provenance area for the Lower Cretaceous Devil's Hole Sandstones to  
 369 its west (Milton-Worrrell *et al.* 2006). These scattered deposits are considered similar to the  
 370 Norwegian Ran Sandstones (Isaksen & Tonstad 1989) and possibly represent a continuation  
 371 of the Upper Jurassic syn-rift Fulmar/Ula shoreface or shelf deposits (Bisewski 1990;  
 372 Johnson & Lott 1993; Copestake *et al.* 2003, Fig. 2). The UK Flora-Fife Trend area and the  
 373 Danish Inge High contain Ula Fm and Rotliegend units in subcrop below the BCU. These are  
 374 potential source areas for the sandstones in the Danish Sten-1 well (Kern *et al.* 1983), which  
 375 is situated in a Lower Cretaceous depocenter (Fig. 7) and is postulated to be a deep marine  
 376 deposit.

377 The 4 m thick unspecified sandstone layer in well NO 2/1-8 (NPD 2013) represents an  
 378 isolated Ran Sandstone occurrence on the Cod Terrace (Fig. 2, 6). The most probable origin  
 379 would be either the Mandal High or the Cod terrace, where well 7/11-8 encounters the  
 380 Skagerrak Fm. in subcrop below the BCU (NPD 2017), indicating a possible small-scale  
 381 sediment provenance area. Any material originating from the Scandinavian mainland to the  
 382 NE would most likely be caught in the Norwegian-Danish Basin region, where major Lower  
 383 Cretaceous depocenters are situated (Copestake *et al.* 2003, Fig. 1a). Similarly, sediments  
 384 from the Josephine High would first have had to cross the Breiflab Basin depocenters (Fig.  
 385 7). However, the exact nature and provenance of these Ran Sandstones cannot be established  
 386 with the data currently available.

387 The thickest Ran sandstones in the study area occur in well NO 3/7-3 (107 m, Fig. 6) and  
 388 these relatively shallow to open marine sandstones were deposited just in the Søgne Basin  
 389 (Verolles 1982; Boirie & Jeannou 1984), which was tectonically inactive during the Lower  
 390 Cretaceous (Rossland *et al.* 2013, Figs. 5, S5, S6). The adjacent Mandal High and its  
 391 metamorphic basement units were largely exposed during the Early Cretaceous (Verolles *et al.*  
 392 1982; Copestake *et al.* 2003; Rossland *et al.* 2013, Fig. 7, 9) and are the probable source  
 393 for these proximal Ran Sandstones (Verolles 1982). Alternatively, Rossland *et al.* (2013)  
 394 suggest, on the basis of dip directions, that these sandstones are related to a turbidite system  
 395 sourced from the Rynkøbing-Fyn High to the east. It should however be stressed that their  
 396 dip-meter data may be affected by post-sedimentary salt movement associated with the large  
 397 salt dome below the Søgne Basin (Verolles 1982, Figs. 5, S5, S6), or could simply represent a

deviation in transport direction as frequently observed within local submarine fan systems (e.g. Normark *et al.* 1979).

The presence of the thick Ran Sandstones in well NO 3/7-3 (Boirie & Jeannou 1984) indicate promising reservoir development in the area, yet none of the other wells in the vicinity encounter Lower Cretaceous sandstones (NPD 2017, Fig. 9). This is in accordance with the depositional character of the Ran Sandstone units described by Verolles (1982) and Boirie & Jeannou (1984), who suggest that reservoir bodies in the area, although potentially of significant thickness, may have a restricted lateral extent (Figs. 9). Furthermore, the Mandal High area is little studied, potentially harbouring reservoirs in various other stratigraphic intervals (Rossland *et al.* 2013, Fig. 10) and detailed analysis will be required to identify these.

In the east of the study area, the Kira Sandstones and their equivalents in the Amalie-1 and Tabita-1 wells (Fig. 6) probably represent submarine fan or slope deposits (Statoil 1991), associated with erosion at BCU-level and the nearby boundary fault between the Tail End Graben and the Ringkøbing-Fyn High (Bonde *et al.* 1994, Fig. 7). Although no rock samples are available from the Amalie-1 well, the metamorphic clasts in cores from the Tabita-1 well are reported to be similar to the basement rocks on the Ringkøbing-Fyn High and on the Mandal High (well NO 3/7-1) (Bonde *et al.* 1994). Possible supply from the Ringkøbing-Fyn High may have involved submarine erosion of the footwall basement, whereas alternative sediment transport from the Mandal High may have by-passed the NO 3/7-3 well and Amalie-1 well before reaching the Tabita-1 well location (Bonde *et al.* 1994, Fig. 7). The sand-prone intervals in the Svane-1 and Iris-1 wells are possibly correlatable to the Kira Sandstones (Bonde *et al.* 1994; Thorsrud *et al.* 2002), which, if correct, may indicate a regional deep marine fan system (Fig. 7). It should be noted however, that except for the Amalie-1 well, no Lower Cretaceous reservoir-quality sandstones are found. Yet a few localised sandy apron or lobe units may have developed as a continuation of the Jurassic deep marine sandstones in the area (Bonde *et al.* 1994; Nielsen *et al.* 2015). Similar deposits could also have developed in the Gertrud Graben and Feda Graben to the South and SW of the exposed Mandal High (Rossland *et al.* 2013), but there is currently no evidence to support this interpretation and identifying such reservoirs, if present, will be highly challenging.

In contrast to the UK and Danish Central Graben areas, no Latest Ryazanian sandstones appear in wells within the NCG proper (NPD 2017) and most Lower Cretaceous depocenters are isolated from the identified sand source areas (Fig. 7). However, various faults were still active, of which some could have exposed sand-prone lithologies to erosion. Of these, the Hydra High block next to the Breiflab Basin, where Rotliegend units are present in the footwall, is the best example (Figs. 5b, 7). However, it is possible that such smaller sand source areas (e.g. Argyll Field area: 10–100 km<sup>2</sup> and less for exposed fault scarps) might not have produced enough sand-prone material for reservoir-size deposits (*sensu* McArthur *et al.* 2016a). By contrast, the exposed Mandal High amounts to 500–600 km<sup>2</sup> and is associated with the thick Ran sandstones in well NO 3/7-3 and the postulated Amalie fan system, thus representing significant reservoir potential.

#### Aptian-Albian reservoir distribution

The interpretation of the Latest Ryazanian reservoir interval is presented in Fig. 8 and depicts the sandstone occurrences in wells, potential source areas with sand-prone lithologies subcropping the Top Lower Cretaceous and sediment transport fairways to depocenters identified on the Lower Cretaceous isochron map.

Towards the end of the Early Cretaceous, sandstone occurrences are rarely seen in UKCG wells (Milton-Worssel *et al.* 2006). However, important sediment provenance areas (e.g. the Forties-Montrose High, Auk Ridge, Josephine High) were still in place and exposed, providing sand influx into the adjacent depocenter as recorded in some penetrations (Milton-Worssel *et al.* 2006, Fig. 8). However, several of the smaller source areas were flooded and covered with Lower Cretaceous deposits (Cod Terrace and Inge High) and potential sourcing from fault scarps was strongly diminished with the cessation of rift activity. Other Ryazanian provenance areas were reduced but remained partially exposed towards the end of the Early Cretaceous as indicated by subcrop data (e.g. the Argyll Field area, Flora-Fife Trend, compare Fig. 8 with Fig. 7), yet no sandstone well occurrences are recorded in the adjacent Aptian-Albian depocenters.

Ran Sandstone units belonging to the Aptian-Albian reservoir interval are found in only one Norwegian well: NO 2/7-15 (Isaksen & Tonstad 1989, NPD 2017, Figs. 6, 8). These clay-rich silt/sandstones are somewhat isolated from the interpreted sediment provenance areas. The Flora Field area, where the Rotliegend is found in subcrop below the Upper Cretaceous chalk deposits, is proposed as the most likely origin of these units (Fig. 8). However, the character of NO 2/7-15 Ran Sandstones remains poorly constrained and demands further assessment.

It should be noted that the wells in the Søgne Basin area, where thick Ryazanian Sandstones were previously deposited (well NO 3/7-3), record only mudstone and chalky deposits (Rossland *et al.* 2013; NPD 2017). Also, the potential Amalie fan system in the Danish Tail End Graben to the south is absent in well reports. Yet the Mandal High was still prone to erosion during the Aptian-Albian, as indicated by metamorphic basement and Bryne Fm subcropping the Upper Cretaceous chalk units (wells NO 2/6-5, NO 3/7-1 and West-Lulu 4, Mærsk 1987; NPD 2017). In addition, large parts of the Ringkøbing-Fyn High have no or thin (a few meters) Lower Cretaceous cover (Japsen *et al.* 2003). Both highs may, therefore, have produced sand-prone material leading to localised reservoir development (Fig. 8), although there is currently no evidence to support this suggestion.

Overall, the Aptian-Albian reservoir interval provides significantly less potential for Lower Cretaceous sandstone deposits than the Latest Ryazanian, due to the drowning of sand source areas. Still, the Ran Sandstone present in well NO 2/7-15 and the sandstone occurrences in various other wells in the UKCG indicate some reservoir potential.

## POTENTIAL ANALOGUE SETTINGS IN THE SOUTHERN PERMIAN BASIN AREA

This study shows that the UKCG and NCG harbour potential for Lower Cretaceous sandstone reservoir units suggesting further exploration possibilities. As the Central Graben structure continues south into Danish, German and Dutch territorial waters (Figs. 1, 11), where the geological setting was quite similar during the Early Cretaceous (Voigt *et al.* 2008; Pharaoh *et al.* 2010), it would be worthwhile to extend the scope of a future case study to these areas.

In Denmark for example, the Ringkøbing-Fyn High along the Eastern margin of the Tail End Graben has no or limited Upper Jurassic to Lower Cretaceous sedimentary cover (Japsen *et al.* 2003), and is known to have been the source of various Late Jurassic fan deposits (Johannessen & Andsbjerg 1993; Andsbjerg & Dybkjær 2003). Such conditions are likely to have continued into at least the Earliest Cretaceous, as illustrated by the deposition of Vyl sandstones (Figs. 2, 11a). These submarine fan units with moderate reservoir potential are found adjacent to the Coffee Soil Fault and were supplied by the Ringkøbing-Fyn High (Michelsen *et al.* 2003, Fig. 11a). In addition, the Lower Cretaceous chalks of the Tuxen Fm. form the reservoirs in the Danish Valdemar and Adda fields (Copestake *et al.* 2003; Jakobsen *et al.* 2005, Fig. 2) indicating another attractive target for continued exploration in the area.

Further to the south, the German and Dutch sectors of the Central Graben are flanked by the Schill Grund High to the east and the Step Graben and Cleaver Bank High to the west (Fig. 11a), areas which were exposed highs during the Late Jurassic and the Early Cretaceous (Pharaoh *et al.* 2010). However, intense Late Cretaceous and Cenozoic basin inversion has caused significant erosion (De Jager 2007) and most of the Lower Cretaceous in the southern sector of the Dutch Central Graben was removed. In the northern sector of the Dutch Central Graben, where inversion and associated erosion was less drastic (Dronkers & Mrozek 1991), Lower Cretaceous sediments are better preserved and hydrocarbon-bearing Scruff sandstones are found (De Jager 2003; De Jager & Geluk 2007, Fig. 2). Additionally, the adjacent Terschelling Basin, where moderate inversion is recorded (Verweij & Witmans 2009) contains relatively thick Lower Cretaceous deposits (Duin *et al.* 2006; EBN *et al.* 2015).

On the southern fringes of the Southern Permian Basin, the Broad Fourteens Basin and West Netherlands Basin form a continuation of the Lower Cretaceous North Sea basins (Fig 11a). Although these basins also underwent strong post-rift inversion (Van Wijhe 1987; De Jager 2003), significant parts of the Lower Cretaceous deposits are preserved in the area (over 900 m thick locally, Duin *et al.* 2006) and contain various hydrocarbon fields (De Jager & Geluk 2007). Similar to the situation in the Moray Firth, the associated reservoirs are documented to be visible on seismic due to a relatively thin Upper Cretaceous-Cenozoic overburden (Oakman 2005). The Early Cretaceous depositional environment was, however, rather different from the situation in the Central Graben and Moray Firth. Instead of isolated shale-dominated basins, receiving limited sand influx from small exposed highs nearby, the area received ample sediment input from the large London-Brabant Massif to the south (Jeremiah *et al.* 2010, Fig. 11a). Therefore, extensive continental to shallow marine shelf clastics were deposited in relatively shallow basins, in contrast with the deep marine basin settings in the Central and Northern North Sea (Figs. 2, 11). The abundance of sand-prone material in the

depositional systems on the fringe of the Southern Permian Basin could potentially have fed submarine fans in the rift depocenters further north. The area experienced tectonically-induced rejuvenation of clastic input, progradation and the development of a widespread shelf system at the K30 sequence boundary, which is associated with increased Hauterivian deep marine reservoir development in the Moray Firth (DeVault & Jeremiah 2002). However, except for the Lower Barremian (Wanneperveen) turbidite units found in association with the Friesland Platform near the Dutch-German border (Jeremiah *et al.* 2010, Fig. 11a), no such deposits are recorded in the Southern Permian Basin. This scarcity of deep marine sand development may be related to the area's relatively gentle bathymetry during the Early Cretaceous (Fig. 11a) although various other factors are known to affect turbidite systems such as shelf width, surrounding geomorphology and hinterland lithologies (Martinsen *et al.* 2005; Mudge 2014).

#### ANALOGUE SETTINGS IN THE SOUTH VIKING GRABEN AREA

Another potential analogue region to the NCG is the South Viking Graben (SVG, Fig. 11). In contrast to the Southern North Sea, the area was associated with a deep marine setting (flanked by exposed highs) during the Earliest Cretaceous (Fig. 11a). Shallow marine or terrestrial sandstones were deposited on the Utsira High, forming parts of the reservoirs in the Edvard Grieg and Johan Sverdrup fields (NPD 2017) and may be directly comparable to the Mandal High in the NCG (Rossland *et al.* 2013). The SVG is documented to include Upper Jurassic turbidites (Partington *et al.* 1993; Fraser *et al.* 2003; Jackson *et al.* 2011). The associated Fladen Ground Spur, Crawford Spur and Utsira High sand provenance areas continued to be exposed in the earliest Cretaceous (Copestake *et al.* 2003, Fig. 11a). However, no Earliest Ryazanian deep marine sands are reported from the SVG area, potentially providing exploration opportunities.

The situation was different during the Aptian, where Skiff Sandstone units are reported along the fringes of the Fladen Ground Spur and the Crawford Spur (Johnson & Lot 1993; Johnson *et al.* 2005, Fig. 11b). To the south, the Kopervik fairway supplied the reservoirs of the giant Britannia Field with sands derived from the East Orkney High in the west (Jeremiah 2000). Oakman (2005) suggests that these deep marine sands represent a fundamentally different depositional system for the Aptian-Albian interval, rather similar to the Cenozoic situation and involving sediment transport over long distances sourced by the exposed North Atlantic rift shoulders, in contrast to the preceding confined Upper Jurassic turbidite fans. The Kopervik system is, however, separated from the SVG by a halokinetically-induced high that was in place throughout the Early Cretaceous, so that potential sandstone deposits in the SVG can only be derived from the adjacent highs (Bisewski 1990, Fig. 11). Further to the north, in the North Viking graben, deep marine slumps of Albian age form reservoirs of the Agat field (Skibeli *et al.* 1995) but these deposits were derived from the main Scandinavian massif (Gulbrandsen & Nyborkken 1991), whereas the SVG remained relatively isolated.

Other wellbore calibrated sandstone occurrences in the area are reported from the Åsta Graben, SE of the Utsira High (3-6 in Fig. 11b, Table 1). These Ran Sandstone units all occur

in the uppermost part of the Lower Cretaceous, directly underneath the Upper Cretaceous chalk deposits and were likely deposited in a shallow marine environment (Olsen 1979; Isaksen & Tonstad 1989).

## POTENTIAL METHODS FOR FURTHER DETAILED RESERVOIR INTERPRETATION

As demonstrated by Milton-Worsell *et al.* (2006), detailed seismic analysis is required to distinguish potential reservoir units. However, the presence of thick Upper Cretaceous chalk and the low impedance contrasts between Cromer Knoll shales and sandstones renders seismic imaging of any relatively thin (typically less than 30 m) Lower Cretaceous sandstone reservoir problematic. To do so requires good quality 3D seismic data combined with an understanding of the likely depositional systems to be encountered (Crittenden *et al.* 1998; Law *et al.* 2000; McKie *et al.* 2015). With such data available, sedimentary systems such as deep marine fans may be traceable on time slice amplitude maps (e.g. Posamentier & Kolla 2003; Martinsen *et al.* 2005; Kilhams *et al.* 2011; 2014a). Amplitude versus offset (AVO) techniques could help to distinguish differences in lithology and reservoir fluid content (e.g. Oakman 2005; Veeken & Rauch-Davies 2006; Milton-Worsell *et al.* 2008; Othman *et al.* 2017). Such a study would be recommended for the Tail End Graben area, as there is potential for small-scale reservoir development. Furthermore, the seismic response of Lower Cretaceous sandstone well occurrences in the NCG, as well as the UKCG where sandstones are more common (Milton-Worsell *et al.* 2006), should be compared to seismic facies in undrilled depocenters. Detailed seismic sequence stratigraphy of Lower Cretaceous depocenters could allow the identification of sea-level driven erosional unconformities on highs, associated with lowstand fans systems in basinal areas (*sensu* Posamentier & Vail 1988).

Methods to further assess sand source areas and to localise associated shallow to deep marine sandstones might include palynological (or similar biostratigraphic) analysis of cored wells to determine to what degree a high was exposed (e.g. O'Driscoll *et al.* 1990; Mudge & Jones 2004; McArthur *et al.* 2016a). Since cores from wells NO 2/7-15 and NO 3/7-3 are available (NPD 2017), magnetic analysis could provide sediment transport directions of these specific Early Cretaceous sandstone occurrences (Hailwood & Ding 2000). Additional petrological and geochemical analysis of heavy minerals (e.g. garnets or zircons) might reveal their provenance area (e.g. Morton *et al.* 2005; Kilhams *et al.* 2014b; Nielsen *et al.* 2015), if cuttings/cores of nearby sand source areas are available (e.g. well NO 3/7-1 on the Mandal High and wells NO 3/7-3 and Tabita-1 in the Søgne Basin and Tail End Graben, respectively; Verolles 1982; Bonde *et al.* 1994, Fig. 7). Furthermore, it will be important to consider the factors influencing the behaviour and geometries of shoreface systems and deep marine fans (e.g. sand-to-mud ratio, flow discharge, slope gradient, sea level changes and fault activity) and where sand deposits occur in these systems (e.g. Posamentier & Kolla 2003; Martinsen *et al.* 2005; McKie *et al.* 2015; McArthur *et al.* 2016b). Recently developed software for the simulation of turbidite deposition in combination with paleorelief reconstructions on 3D seismic could be a powerful tool to predict the distribution of deep marine fans (Aas *et al.* 2010).

608

## CONCLUSIONS

609 Here a structural framework of the NCG area has been presented. This reflects a diverse  
610 geological history including Triassic extension and salt movement, Late Jurassic to Early  
611 Cretaceous rifting and subsequent basin inversion with salt diapirism. Late Jurassic rifting  
612 was most intense in the south of the study area, while Early Cretaceous rifting was more  
613 important in the north, possibly representing an Early Cretaceous change in tectonic regime  
614 before rifting halted altogether. An assessment of the Lower Cretaceous indicates fair  
615 potential for reservoir development. Although the study area is isolated from a large  
616 hinterland, local sediment sources and potential sediment transport routes are identified. Most  
617 potential is expected around the exposed highs in Ryazanian times, while many sand source  
618 areas were drowned at the end of the Early Cretaceous (Aptian-Albian). The underexplored  
619 Mandal-High area, where restricted shallow marine sandstone deposits around the exposed  
620 Mandal High and in the Søgne Basin provide the best potential, is suggested for further focus.  
621 Similar depositional environments could have existed around other exposed highs (e.g.  
622 Josephine High, Auk Ridge), although they may have been too small to have produced  
623 significant reservoir units. Furthermore, the postulated Amalie fan system in Denmark  
624 illustrates the possibilities for good quality deep marine sandstones, which may have also  
625 formed in the depocenters south and SW of the Mandal High. Analogous settings to those in  
626 the study area are also recognised in the Southern Permian Basin area to the south and the  
627 South Viking Graben to the north, further analysis of the Lower Cretaceous reservoir  
628 intervals of these areas would be an interesting next step. A detailed effort including the use  
629 of advanced seismic techniques and detailed well analysis will be necessary to accurately  
630 define such reservoirs, if present. The discovery of the Edvard Grieg and Johan Sverdrup  
631 fields illustrates the importance of continued exploration, especially the re-assessment of  
632 available well and seismic data, in the context of this mature hydrocarbon province (Jørstad  
633 2012).

634

635

636 This project was originally undertaken as an internship at Shell Upstream International in  
637 Assen, Netherlands as the basis for a master's project (VU University Amsterdam). I would  
638 first like to express my gratitude to Carlo Nicolai and Ben Kilhams (Royal Dutch Shell) as  
639 well as Jan de Jager (VU University Amsterdam) for their support and guidance during this  
640 project and for helping me to prepare this manuscript for publication. I would also like to  
641 thank my colleagues from the Shell's UK/Netherlands Exploration Teams, the NAM IT team  
642 and colleagues from Shell Norway for their help and support. I am grateful to Adam  
643 McArthur and Jason Jeremiah for their detailed and constructive comments and to Royal  
644 Dutch Shell for funding this project and for allowing me to publish its results.

645



## References

- A/S Norske Shell. 1968. Exploration Well Resumé. Well 17/11-1. A/S Norske Shell Exploration & Petroleum Engineering Departments.  
[http://www.npd.no/engelsk/cwi/pbl/wellbore\\_documents/906\\_01\\_Exploration\\_well\\_resume.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/906_01_Exploration_well_resume.pdf)
- Aas, T.E., Howell, J.A., Janocko, M. & Jackson, C.A.-L. 2010. Control of Aptian palaeobathymetry on turbidite distribution in the Buchan Graben, Outer Moray Firth, Central North Sea. *Marine and Petroleum Geology*, **27**, 412-434.  
 doi:10.1016/j.marpetgeo.2009.10.014
- Ainsworth, N.R., Riley, L.A. & Gallagher, L.T. 2000. An Early Cretaceous lithostratigraphic and biostratigraphic framework for the Britannia Field reservoir (Late Barremian–Late Aptian), UK North Sea. *Petroleum Geoscience*, **6**, 345–367.  
 DOI: 10.1144/petgeo.6.4.345
- Andsbjerg, J. & Dybkjær, K. 2003 Sequence stratigraphy of the Jurassic of the Danish Central Graben. *In: Ineson, J.R. & Surlyk, F. (eds) The Jurassic of Denmark and Greenland. Geological Survey of Denmark and Greenland Bulletin*, **1**, *Geological Survey of Denmark and Greenland Bulletin*, **1**, 265–300.
- Argent, J.D., Stewart, S.A. & Underhill, J.R. 2000. Controls on the Lower Cretaceous Punt Sandstone Member, a massive deep-water clastic deposystem, Inner Moray Firth, UK North Sea. *Petroleum Geoscience*, **6**, 275-285. <https://doi.org/10.1144/petgeo.6.3.275>
- Bartholomew, I.D., Peters, J.M. & Powell, C.M. 1993. Regional structural evolution of the North Sea: oblique slip and the reactivation of basement lineaments. *In: Parker, J.R. (ed) Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 4th Petroleum Geology Conference*. . Geological Society, London, Petroleum Geology Conference series, **4**, 1109–1122.  
 DOI: 10.1144/0041109
- Bergan, M., Tørudbakken, B. & Wandås, B. 1989. Lithostratigraphic correlation of Upper Jurassic sandstones within the Norwegian Central Graben: sedimentological and tectonic implications. *In: Collinson, J. (ed) Correlation in Hydrocarbon Exploration*. Graham & Trotman, London, 243–251. DOI: 10.1007/978-94-009-1149-9\_19
- Bisewski, H. 1990. Occurrence and depositional environment of the Lower Cretaceous sands in the southern Witch Ground Graben. *In: Hardman, R.F.P. & Brooks, J. (eds), Tectonic Events Responsible for Britain's Oil and Gas Reserves*. Geological Society Special Publication, **55**, 325-338.

- Blackbourn, G.A. & Thomson, M.E. 2000. Brittania Field, UK North Sea: petrographic constraints on Cretaceous provenance, facies and the origin of slurry-flow deposits. *Petroleum Geoscience*, **6**, 329-343.
- Bonde, K., Sandvik, R., Slot-Petersen, C. & Olsen, T. 1994. Tabita-1, Tabita-1A Completion Report, Statoil Efterforskning OG Produktion A/S, Copenhagen, Denmark.
- Boirie, J.M. & Jeannou, A. 1984. Well 3/7-3 Lithostratigraphy and reservoir aspect of the pre-chalk sediments (Permian – Lower Cretaceous; Central Graben) Southern District – Offshore Norway. Boussens well report, **062/84 RP**.  
[http://www.npd.no/engelsk/cwi/pbl/wellbore\\_documents/293\\_03\\_3\\_7\\_3\\_LITHOSTRATIGRAPHY\\_AND\\_RESERVOIR\\_ASPECT\\_OF\\_THE\\_PRE-CHALK\\_SEDIMENTS.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/293_03_3_7_3_LITHOSTRATIGRAPHY_AND_RESERVOIR_ASPECT_OF_THE_PRE-CHALK_SEDIMENTS.pdf)
- Britoil. 1985. Iris-1 composite well log. Britoil PLC. Exploration drawing office, Glasgow, UK.
- Brooks, J. 1990. Classic petroleum provinces. In: Brooks, J. (ed) *Classic Petroleum Provinces*. Geological Society, London, Special Publications, **50**, 1–8. DOI: 10.1144/GSL.SP.1990.050.01.01
- Cartwright, J.A. 1989. The kinematics of inversion in the Danish Central Graben. In: Cooper, M. A., & Williams, G. D. (eds) *Inversion Tectonics*. Geological Society, London, Special Publications, **44**, 153–175. DOI: 10.1144/GSL.SP.1989.044.01.10
- Copestake, P., Sims, A.P., Crittenden, S., Hamar, G.P., Ineson, J.R., Rose, P.T. & Tringham, M.E. 2003. Lower Cretaceous. In: Evans, D., Graham, C., Armour, A. & Bathurst, P. (eds) *The Millennium Atlas: Petroleum Geology of the Central and Northern North Sea*. Geological Society, London, 191–211.
- Coward, M.P., Dewey, J.F., Hempton, M. & Holroyd, J. 2003. Tectonic evolution. In: Evans, D., Graham, C., Armour, A. & Bathurst, P. (eds) *The Millennium Atlas: Petroleum Geology of the Central and Northern North Sea*. Geological Society, London, 17-33.
- Crittenden, S., Cole, J.M. & Kirk, M.J. 1997. THE DISTRIBUTION OF APTIAN SANDSTONES IN THE CENTRAL AND NORTHERN NORTH SEA (UK SECTORS): A LOWSTAND SYSTEMS TRACT “PLAY”. Part 1. Stratigraphy, biostratigraphic age determination, and genesis of the sandstones. *Journal of Petroleum Geology*, **20** (1), 3-25.
- Crittenden, S., Cole, J.M. & Kirk, M.J. 1998. THE DISTRIBUTION OF APTIAN SANDSTONES IN THE CENTRAL AND NORTHERN NORTH SEA (UK SECTORS) - A LOWSTAND SYSTEMS TRACT PLAY. Part 2. distribution and exploration strategy. *Journal of Petroleum Geology*, **21** (2), 187-211.
- De Jager, J. 2003. Inverted Basins in the Netherlands, similarities and differences. *Netherlands Journal of Geosciences/Geologie en Mijnbouw*, **82**, 4, 355–366.

doi: <https://doi.org/10.1017/S0016774600020175>

De Jager, J. 2007. Geological Development. *In*: Wong, Th.E., Batjes, D.A.J. & De Jager, J. (eds) *Geology of the Netherlands*. Royal Netherlands Academy of Science and Art, Amsterdam, 5–26.

De Jager., J. & Geluk, M.C. 2007. Petroleum Geology. *In*: Wong, Th.E., Batjes, D.A.J. & De Jager, J. (eds) *Geology of the Netherlands*. Royal Netherlands Academy of Science and Art, Amsterdam, 241–264.

DeVault, B. & Jeremiah, J. 2002. Tectonostratigraphy of the Nieuwerkerk Formation (Delfland subgroup), West Netherlands Basin. *AAPG Bulletin*, **86** (10), 1679-1707. DOI: 10.1306/61EEDD50-173E-11D7-8645000102C1865D

Dronkers, A.J. & Mrozek, F.J. 1991. Inverted basins of The Netherlands. *First Break*, **9**, 409–425. DOI: 10.3997/1365-2397.1991019

Duin, E.J.T., Doornebal, J.C., Rijkers, R.H., Verbeek, J.W. & Wong, Th. E. 2006. Subsurface structure of the Netherlands – results of recent onshore and offshore mapping. *Netherlands Journal of Geosciences/Geologie en Mijnbouw*, **85** (4), 355–366.

EBN, Ministry of Economic Affairs & TNO, 2015. Prospectivity G and M blocks – Analysing the Upper Jurassic play. <http://www.nlog.nl/cm15/browser?id=workspace%3A//SpacesStore/f84ec4ac-b9a6-4771-9212-6f6959233ead>

Eggenhuisen, J.T., McCaffrey, W.D., Haughton, P.D.W., Butler, R.W.H., Moore, I., Jarvie, A., Hakes, W.G. 2010. Reconstructing large-scale remobilisation of deep-water deposits and its impact on sand-body architecture from cored wells: The Lower Cretaceous Britannia Sandstone Formation, UK North Sea. *Marine and Petroleum Geology*, **27**, 1595-1615. doi:10.1016/j.marpetgeo.2010.04.005

Erratt, D., Thomas, G.M. & Wall, G.R.T. 1999. The evolution of the Central North Sea Rift. Geological Society, *In*: Fleet, A.J. & Boldy, S.A.R. (eds) *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 5th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **5**, 63–82. DOI: 10.1144/0050063

Fjellanger, E. 1986. Geological Completion Report, Well 2/1-8. BP Petroleum Development (Norway) Ltd. U/A, Stavanger, Norway. [http://www.npd.no/engelsk/cwi/pbl/wellbore\\_documents/459\\_2\\_1\\_8\\_Completion\\_report.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/459_2_1_8_Completion_report.pdf)

Fraser, S.I., Robinson, A.M., Johnson, H.D., Underhill, J.R. & Kadolsky, D.G.A., Connell, R., Johannessen, P. & Ravnås, R. 2003. Upper Jurassic. *In*: Evans, D., Graham, C.,

Armour, A. & Bathurst, P. (eds) *The Millennium Atlas: Petroleum Geology of the Central and Northern North Sea*. Geological Society, London, 155–189.

Fossen, H., Dallman, W. & Andersen, T.B. 2008. The mountain chain rebounds and founders. In: Ramberg, I.B., Bryhni, I., Nøttvedt, A. & Rangnes, K. (eds) *The Making of a Land - Geology of Norway*. Norsk Geologisk Forening. 232–259.

Garrett, S.W., Atherton, T. & Hurst, A. 2000. Lower Cretaceous deep-water sandstone reservoirs in the UK Central North Sea. *Petroleum Geoscience*, **6**, 231–240.  
<https://doi.org/10.1144/petgeo.6.3.231>

Gautier, D.L. 2005. Kimmeridge Shales Total Petroleum System of the North Sea Graben Province. *U.S. Geological Survey Bulletin*, **2204-C**.  
<https://pubs.usgs.gov/bul/2204/c/pdf/B2204C.pdf>

Ge, Z., Gawthorpe, R.L., Rotevatn, A. & Thomas, M.B. 2016. Impact of normal faulting and pre-rift salt tectonics on the structural style of salt-influenced rifts: the Late Jurassic Norwegian Central Graben, North Sea. *Basin Research*, 1–25. doi: 10.1111/bre.12219

Gowers, M.B. & Sæbøe, A. 1985. On the structural evolution of the Central Trough in the Norwegian and Danish sectors of the North Sea. *Marine and Petroleum Geology*, **2**, 298–318. DOI:10.1016/0264-8172(85)90026-1

Gowers, M.B., Holtar, E. & Swensson, E. 1993. The structure of the Norwegian Central Trough (Central Graben area). In: Parker, J.R. (ed) *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 4th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **4**, 1245–1254. DOI: 10.1144/0041245

Graversen, O. 2006. The Jurassic-Cretaceous North Sea Rift Dome and Associated Basin Evolution. *AAPG Search and Discovery Article*, **#3004**.  
<http://www.searchanddiscovery.com/documents/2006/06013graversen/images/graversen>

Gulbrandsen, A. & Nyborkken, S. 1991. Agat Field - Norwegian North Sea, Subplatform North Viking Graben. *AAPG Search and Discovery Article*, **#91004**.  
<http://www.searchanddiscovery.com/abstracts/html/1991/annual/abstracts/0586b.htm>

Hampson, J.M., Walden, S.F. & Bell, C. 2010. Maximizing production and reserves from offshore heavy oil fields using seismic and drilling technology: Alba and Captain Fields, UKNS. In: Vining, B.A. & Pickering, S.C. (eds) *Petroleum Geology: From Mature Basins to New Frontiers – Proceedings of the 7th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **7**, 337–347.  
DOI: 10.1144/0070337

- 822 Hailwood, E. & Ding, F. 2000. Sediment transport and dispersal pathways in the Lower  
823 Cretaceous sands of the Britannia Field, derived from magnetic anisotropy. *Petroleum*  
824 *Geology*, **6**, 369-379. <https://doi.org/10.1144/petgeo.6.4.369>  
825
- 826 Herngreen, G.F.W. & Wong, Th.E. 2007. Cretaceous. *In*: Wong, Th.E., Batjes, D.A.J. &  
827 De Jager, J. (eds) *Geology of the Netherlands*. Royal Netherlands Academy of Science and  
828 Art, Amsterdam, 127–150.  
829
- 830 Hodgson, N.A., Fansworth, J. & Fraser, A.J. 1992. Salt-related tectonics, sedimentation  
831 and hydrocarbon plays in the Central Graben, North Sea, UKCS. *In*: Hardman, R.F.P. (ed)  
832 *Exploration Britain: Geological insights for the next decade*. Geological Society, London,  
833 Special Publications, **67**, 31–63. DOI: 10.1144/GSL.SP.1992.067.01.03  
834
- 835 Isaksen, D. & Tonstad, K. 1989. A revised Cretaceous and Tertiary lithostratigraphic  
836 nomenclature for the Norwegian North Sea. *Norwegian Petroleum Directorate Bulletin*, **5**.  
837 <http://www.npd.no/Global/Norsk/3-Publikasjoner/NPD-Bulletin/Bulletinnr5.pdf>  
838
- 839 Jakobsen, F., Ineson, J.R., Kirstensen, L., Nytoft, H.P. & Stemmerik, L. 2005. The  
840 Valdemar Field, Danish Central Graben: field compartmentalization and regional  
841 prospectivity of the Lower Cretaceous chalk play. *In*: Doré, A.G. & Vining, B.A. (eds)  
842 *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 6th*  
843 *Petroleum Geology Conference*. Geological Society, London, Petroleum Geology  
844 Conference series, 6, 177-186. <https://doi.org/10.1144/0060177>  
845
- 846 Jackson, C.A.-L., Larsen, E., Hanslien, S., Tjemsland, A.-E. 2011. Controls on synrift  
847 turbidite deposition on the hanging wall of the South Viking Graben, North Sea rift  
848 system, offshore Norway. *AAPG Bulletin*, **95** (9), 1557-1587. DOI:10.1306/01031110037  
849
- 850 Jackson, C.A.-L. & Lewis, M.M. 2016. Structural style and evolution of a salt-influenced  
851 rift basin margin; the impact of variations in salt composition and the role of polyphase  
852 extension. *Basin Research*, **28**, 81-102. doi: 10.1111/bre.12099  
853
- 854 Japsen, P., Britze, P. & Andersen, C. 2003. Upper Jurassic - Lower Cretaceous of the  
855 Danish Central Graben: structural framework and nomenclature. *In*: Ineson, J.R. & Surlyk,  
856 F. (eds) *The Jurassic of Denmark and Greenland*. Geological Survey of Denmark and  
857 Greenland Bulletin, **1**, 233–246. [http://www.geus.dk/DK/publications/geol-survey-dk-gl-](http://www.geus.dk/DK/publications/geol-survey-dk-gl-bull/Documents/nr1_p231-246.pdf)  
858 [bull/Documents/nr1\\_p231-246.pdf](http://www.geus.dk/DK/publications/geol-survey-dk-gl-bull/Documents/nr1_p231-246.pdf)  
859
- 860 Jeremiah, J.M. 2000. Lower Cretaceous turbidites of the Moray Firth: sequence  
861 stratigraphical framework and reservoir distribution. *Petroleum Geoscience*, **6**, 309–328.  
862 DOI: 10.1144/petgeo.6.4.309  
863
- 864 Jeremiah, J.M., Duxbury, S. & Rawson, P. 2010. Lower Cretaceous of the southern North  
865 Sea Basins: reservoir distribution within a sequence stratigraphic framework. *Netherlands*  
866 *Journal of Geosciences – Geologie en Mijnbouw*, **89** (3/4). 203-237.

Doi: <https://doi.org/10.1017/S0016774600000706>

Johannessen, P.N. & Andsbjerg, J. 1993. Middle to Late Jurassic basin evolution and sandstone reservoir distribution in the Danish Central Trough. *In: Parker, J.R. (ed) Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 4th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **4**, 271–283. doi: 10.1144/0040271

Johnson, J. & Lott, G.K. 1993. 2. Cretaceous of the Central and Northern North Sea. *In: Knox, R.W.O'B. & Cordey, W.G. (eds) Lithostratigraphic nomenclature of the UK North Sea*. British Geological Survey (Nottingham).  
<http://www.bgs.ac.uk/downloads/start.cfm?id=2044>

Johnson, H., Leslie, A.B., Wilson, C.K., Andrews, I.J. & Cooper, R.M. 2005. Middle Jurassic, Upper Jurassic and Lower Cretaceous of the UK Central and Northern North Sea. *British Geological Survey Research Report*, **RR/03/001**.  
<http://nora.nerc.ac.uk/3680/1/RR03001.pdf>

Jones, L.S., Garrett, S.W., MacLeod, M., Guy, M., Codon, P.J. & Notman, L. 1999. Britannia Field, UK Central North Sea: modelling heterogeneity in unusual deep-water deposits. *In: Fleet, A.J. & Boldy, S.A.R. (eds) Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 5th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **5**, 1115-1124.  
<https://doi.org/10.1144/0051115>

Jørstad, A.J. 2012. Johan Sverdrup – Offshore Norway: The Story behind the Giant Sverdrup Discovery. *AAPG Search and Discovery Article*, **#20177**.  
[https://www.google.ch/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=0ahUKEwiL4v6vwurVAhXGJMAKHxveBpQQFggyMAA&url=http%3A%2F%2Fwww.searchanddiscovery.com%2Fdocuments%2F2012%2F20177jorstad%2Fndx\\_jorstad.pdf&usg=AFQjCNErkXDHLyXjqcp8dlbV\\_ZJuqMtKYQ](https://www.google.ch/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=0ahUKEwiL4v6vwurVAhXGJMAKHxveBpQQFggyMAA&url=http%3A%2F%2Fwww.searchanddiscovery.com%2Fdocuments%2F2012%2F20177jorstad%2Fndx_jorstad.pdf&usg=AFQjCNErkXDHLyXjqcp8dlbV_ZJuqMtKYQ)

Kern, J.C., Rørvig, M., Liboriussen, J., Blom, N.S. & Muelder, J. 1983. Sten-1 completion report. Chevron Petroleum Company of Denmark, Copenhagen, Denmark.

Kilhams, B., Godfrey, S., Harlley, A. & Huuse, M. 2011. An integrated 3D seismic, petrophysical and analogue core study of the Mid-Eocene Grid channel complex in the greater Nelson Field area, UK Central North Sea. *Petroleum Geoscience*, **17**, 127-142. DOI 10.1144/1354-079310-022

Kilhams, B., Hartley, A., Huuse, M. & Davis, C. 2014a. Characterizing the Paleocene turbidites of the North Sea: Maureen Formation, UK Central Graben. *In: McKie, T., Rose, P.T.S., Hartley, A.J., Jones, D.W. & Armstrong, T.L. (eds) Tertiary Deep-Marine Reservoirs of the North Sea Region*. Geological Society, London, Special Publications, **403**, 43-62.

<http://dx.doi.org/10.1144/SP403.1>

Kilhams, B., Morton, A., Borella, R., Wilkins, A., Hurst, A. 2014b. Understanding the provenance and reservoir quality of the Sele Formation sandstones of the UK Central Graben utilizing detrital garnet suites. *In: Scott, R.A., Smyth, H.R., Morton, A.C. & Richardson, N. (eds) 2014. Sediment Provenance Studies in Hydrocarbon Exploration and Production*. Geological Society, London, Special Publications, **386**, 129–142. <https://doi.org/10.1144/SP386.16>

Law, A., Raymond, A., White, G., Atkinson, A., Clifton, M., Atherton, T., Dawes, I.a.n., Robertson, E., Melvin, A & Brayley, S. 2000. The Kopervik fairway, Moray Firth, UK. *Petroleum Geoscience*, **6**, 265–274. DOI: 10.1144/petgeo.6.3.265

Lundin. 2017. Johan Sverdrup Development. *Lundin Petroleum press releases*. [https://www.lundin-petroleum.com/download/ot\\_johansverdrup\\_present\\_e/?wpdmdl=6265/](https://www.lundin-petroleum.com/download/ot_johansverdrup_present_e/?wpdmdl=6265/)

Mærsk. 1987. Final Well Report West Lulu-4. Mærsk Olie OG Gas A/S Petroleum Engineering and Operations Department

Martinsen, O.J., Lien, T. & Jackson, C.J. 2005. Cretaceous and Palaeogene turbidite systems in the North Sea and Norwegian Sea Basins: source, staging area and basin physiography controls on reservoir development. *In: Doré, A.G. & Vining, B.A. (eds) Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 6th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **6**, 1147–1164. <https://doi.org/10.1144/0061147>

McArthur, A.D., Jolley, D.W., Hartley, A.J., Archer, S.G. & Lawrence, H.M. 2016a. Palaeoecology of syn-rift topography: A Late Jurassic footwall island on the Josephine Ridge, Central Graben, North Sea. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **459**, 63–75. <http://dx.doi.org/10.1016/j.palaeo.2016.06.033>

McArthur, A.D., Hartley, A.J., Archer, S.G., Jolley, D.W. & Lawrence, H.M. 2016b. Spatiotemporal relationships of deep-marine, axial, and transverse depositional systems from the synrift Upper Jurassic of the central North Sea. *AAPG Bulletin*, **100** (9), 1469–1500. DOI:10.1306/04041615125

McGann, G.J., Green, S.C.H., Harker, S.D. & Romani, R.S. 1991. The Scapa Field, Block 14/19, UK North Sea. *In: Abbotts, I.L. (ed). United Kingdom Oil and Gas Fields, 25 Years Commemorative Volume*. Geological Society, London, Memoirs, **14**, 369–376. doi: 10.1144/GSL.MEM.1991.014.01.46

McKie, T., Rose, P.T.S., Hartley, A.J., Jones, D.W. & Armstrong, T.L. 2015. Tertiary deep-marine reservoirs of the North Sea region: an introduction. *In: McKie, T., Rose, P.T.S., Hartley, A.J., Jones, D.W. & Armstrong, T.L. (eds) Tertiary Deep-Marine*

*Reservoirs of the North Sea Region*. Geological Society, London, Special Publications, **403**, 1–16. DOI: 10.1144/SP403.12

Michelsen, O., Nielsen, L.H., Johannessen, P.N., Andsbjerg, J. & Surlyk, F. 2003. Jurassic lithostratigraphy and stratigraphic development onshore and offshore Denmark. *In*: Ineson, J.R. & Surlyk, F. (eds) *The Jurassic of Denmark and Greenland*. Geological Survey of Denmark and Greenland Bulletin, **1**, 147–216.  
[http://www.geus.dk/DK/publications/geol-survey-dk-gl-bull/Documents/nr1\\_p145-216.pdf](http://www.geus.dk/DK/publications/geol-survey-dk-gl-bull/Documents/nr1_p145-216.pdf)

Milton, N.J. 1993. Evolving depositional geometries in the North Sea Jurassic rift. *In*: Parker, J.R. (ed) *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 4th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **4**, 425–442. DOI: 10.1144/0040425

Milton-Worssell, R.J., Stoker, S.J. & Cavill, J.E. 2006. Lower Cretaceous deep-water sandstone plays in the UK Central Graben. *In*: Allen, M.R., Goffey, G.P. Morgan, R.K. & Walker, I.M. (eds) *The deliberate Search for the Stratigraphic Trap*. Geological Society, London, Special Publications, **254**, 169–186. DOI: 10.1144/GSL.SP.2006.254.01.09

Milton-Worssell, R., Fisher, G., Cameron, D., Stoker, S., Avu, A. & Basford, H. 2008. WHERE ARE THOSE LOWER CRETACEOUS SANDS? GEOBODY DETECTION IN A DEPOCENTRE. *PETEX conference proceedings*, Petroleum Exploration Society of Great Britain (PESGB), London.  
[https://www.google.ch/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=0ahUKEwiK7q-o8uzVAhXHC8AKHdqICIEQFggwMAA&url=http%3A%2F%2Fwww.geoteric.com%2Fhubfs%2Fuploads%2Fdownloads%2FPETEX\\_2008.pdf%3Ft%3D1501511671995&usg=AFQjCNE4aS36bSsk5ytaP10DBzZaFZ6hWg](https://www.google.ch/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=0ahUKEwiK7q-o8uzVAhXHC8AKHdqICIEQFggwMAA&url=http%3A%2F%2Fwww.geoteric.com%2Fhubfs%2Fuploads%2Fdownloads%2FPETEX_2008.pdf%3Ft%3D1501511671995&usg=AFQjCNE4aS36bSsk5ytaP10DBzZaFZ6hWg)

Møller, J.J. & Rasmussen, E.S. 2003. Middle Jurassic – Early Cretaceous rifting of the Danish Central Graben. *In*: Ineson, J.R. & Surlyk, F. (eds) *The Jurassic of Denmark and Greenland*. Geological Survey of Denmark and Greenland Bulletin, **1**, 247–264.  
[http://www.geus.dk/DK/publications/geol-survey-dk-gl-bull/Documents/nr1\\_p247-264.pdf](http://www.geus.dk/DK/publications/geol-survey-dk-gl-bull/Documents/nr1_p247-264.pdf)

Morton, A.C., Whitman, A.G. & Fanning, C.M. 2005. Provenance of Late Cretaceous to Paleocene submarine fan sandstones in the Norwegian Sea: Integration of heavy mineral, mineral chemical and zircon age data. *Sedimentary Geology*, **182**, 3–28.  
<https://doi.org/10.1016/j.sedgeo.2005.08.007>

Mudge, D.C. & Jones, S.M. 2004. Palaeocene uplift and subsidence events in the Scotland–Shetland and North Sea region and their relationship to the Iceland Plume. *Journal of the Geological Society, London*, **161**, 381–386.  
<https://doi.org/10.1144/0016-764903-038>



- 1002
- 1003 Mudge, D.C. 2014. Regional controls on Lower Tertiary sandstone distribution in the
- 1004 North Sea and NE Atlantic margin basins. *In: McKie, T., Rose, P.T.S., Hartley, A.J.,*
- 1005 *Jones, D.W. & Armstrong, T.L. (eds.) Tertiary Deep-Marine Reservoirs of the North Sea*
- 1006 *Region*. Geological Society, London, Special Publications, **403**.
- 1007 <https://doi.org/10.1144/SP403.5>
- 1008
- 1009 NDD (Norway Discovery Digest). 2012. The HPHT and UHPHT Play in the Norwegian
- 1010 Central Graben. Canadian Discovery Ltd., **13**, 12–33.
- 1011 <http://norwaydigest.canadiandiscovery.com/?q=node/33144>
- 1012
- 1013 Nielsen, M.T., Weibel, R., Friis, H. 2015. Provenance of gravity-flow sandstones from the
- 1014 Upper Jurassic-Lower Cretaceous Farsund Formation, Danish Central Graben, North Sea.
- 1015 *Marine and Petroleum Geology*, **59**, 371-389.
- 1016 <http://dx.doi.org/10.1016/j.marpetgeo.2014.09.016>
- 1017
- 1018 Normark, W.R., Piper, D.J.W. & Hess, G.R. 1979. Distributary channels, sand lobes, and
- 1019 mesotopography of Navy Submarine Fan, California Borderland, with applications to
- 1020 ancient fan sediments. *Sedimentology*, **26**, 749-774. DOI: 10.1111/j.1365-
- 1021 3091.1979.tb00971.x
- 1022
- 1023 Nøttvedt, A. & Johannessen, E.P. 2008. The source of Norway's oil wealth. *In: Ramberg,*
- 1024 *I.B., Bryhni, I., Nøttvedt, A. & Rangnes, K. (eds) The Making of a Land - Geology of*
- 1025 *Norway*. Norsk Geologisk Forening, 384–417
- 1026
- 1027 NPD. 1979a. Interpreted lithology log. Well 7/3-1. *NPD Paper*, **20**.
- 1028 [http://www.npd.no/engelsk/cwi/pbl/NPD\\_papers/164\\_02\\_NPD\\_Paper\\_No.20\\_Interpreted](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/164_02_NPD_Paper_No.20_Interpreted_Lithology_log__Well_7_3_1.pdf)
- 1029 [\\_Lithology\\_log\\_\\_Well\\_7\\_3\\_1.pdf](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/164_02_NPD_Paper_No.20_Interpreted_Lithology_log__Well_7_3_1.pdf)
- 1030
- 1031 NPD. 1979b. Interpreted lithology log. Well 17/10-1. *NPD Paper*, **21**.
- 1032 [http://www.npd.no/engelsk/cwi/pbl/NPD\\_papers/161\\_02\\_NPD\\_Paper\\_No.21\\_Interpreted](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/161_02_NPD_Paper_No.21_Interpreted_Lithology_log__Well_17_10_1.pdf)
- 1033 [\\_Lithology\\_log\\_\\_Well\\_17\\_10\\_1.pdf](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/161_02_NPD_Paper_No.21_Interpreted_Lithology_log__Well_17_10_1.pdf)
- 1034
- 1035 NPD. 2013. Wellbore 2/1-8. Composite log.
- 1036 [http://www.npd.no/engelsk/cwi/pbl/wellbore\\_composite\\_logs/459.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_composite_logs/459.pdf)
- 1037
- 1038 NPD. 2014. Lithostratigraphic Chart Norwegian North Sea. **OD 1409001**.
- 1039 [http://www.npd.no/Global/Engelsk/2-Topics/Geology/Lithostratigraphy/NS-](http://www.npd.no/Global/Engelsk/2-Topics/Geology/Lithostratigraphy/NS-OD1409001.pdf)
- 1040 [OD1409001.pdf](http://www.npd.no/Global/Engelsk/2-Topics/Geology/Lithostratigraphy/NS-OD1409001.pdf)
- 1041
- 1042 NPD. 2017. (Norwegian Petroleum Directorate website and FactPages/Maps)
- 1043 [www.npd.no](http://www.npd.no) and <http://factpages.npd.no/factpages/Default.aspx?culture=en>
- 1044

- Oakman, C. D. & Partington, M. A. 1998. Cretaceous. *In*: Glennie, K. W. (ed) *Petroleum Geology of the North Sea – Basic Concepts and Recent Advances*. 4th edition, Blackwell, Oxford, 294–349.
- Oakman, C.D. 2005. The Lower Cretaceous plays of the Central and Northern North Sea: Atlantean drainage models and enhanced hydrocarbon potential. *In*: Doré, A.G. & Vining, B.A. (eds) *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 6th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **6**, 187–198. DOI: 10.1144/0060187
- O’Driscoll, D., Hindle, A.D., & Long, D.C. 1990. The structural controls on Upper Jurassic and Lower Cretaceous reservoir sandstones in the Witch Ground Graben, UK North Sea. *In*: Hardman, R.F.P. & Brooks, J. (eds) *Tectonic Events Responsible for Britain's Oil and Gas Reserves*. Geological Society Special Publication, **55**, 299-323.
- Olsen, R.C. 1979. Lithology. Well 17/10-1. *NPD Paper*, **21**.  
[http://www.npd.no/engelsk/cwi/pbl/NPD\\_papers/161\\_01\\_NPD\\_Paper\\_No.21\\_Lithology\\_-\\_Well\\_17\\_10\\_1.pdf](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/161_01_NPD_Paper_No.21_Lithology_-_Well_17_10_1.pdf)
- Othman, A.A.A., Bakr, A. & Maher, A. 2017. Integrated seismic tools to delineate Pliocene gas-charged geobody, offshore west Nile delta, Egypt. *NRIAG Journal of Astronomy and Geophysics*, **6**, 81-89. <http://dx.doi.org/10.1016/j.nrjag.2017.04.008>
- Partington, M.A., Mitchener, B.C., Milton, N.J. & Fraser, A.J. 1993. Genetic sequence stratigraphy for the North Sea Late Jurassic and Early Cretaceous: distribution and prediction of Kimmeridgian–Late Ryazanian reservoirs in the North Sea and adjacent areas. *In*: Parker, J.R. (ed) *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 4th Petroleum Geology Conference*. Geological Society, London, Petroleum Geology Conference series, **4**, 347-370. doi: 10.1144/0040347
- Pharaoh, T.C., Duser, M., Geluk, M.C., Kockel, F., Krawczyk, C.M., Krzywiec, P., Scheck-Wenderoth, M., Thybo, H., Vejbæk, O.V. & Van Wees, J.D. 2010. Tectonic Evolution. *In*: Doornenbal, J.C. & Stevenson, A.G. (eds) *Petroleum Geological Atlas of the Southern Permian Basin Area*. EAGE Publications b.v., Houten, 25–58.
- Phillips, 1981. Completion book for the exploratory well 2/7-15. Phillips Petroleum Company Norway.  
[http://www.npd.no/engelsk/cwi/pbl/wellbore\\_documents/225\\_01\\_2\\_7\\_15\\_Completion\\_report.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/225_01_2_7_15_Completion_report.pdf)
- Pinnock, S.J., Clitheroe, A.R.J. & Rose, P.T.S. 2003. The Captain Field, Block 13/22a, UK North Sea. *In*: Gluyas, J.G. & Hitchens, H.M. (eds) *United Kingdom Oil and Gas Fields, Commemorative Volume*. Geological Society, London, *Memoirs*, **20**, 431-441. doi: 10.1144/GSL.MEM.2003.020.01.35

- 1090 Posamentier, H.W. & Vail, P.R. 1988. EUSTATIC CONTROLS ON CLASTIC  
1091 DEPOSITION II—SEQUENCE AND SYSTEMS TRACT MODELS. *In*: Wilgus, C.K.,  
1092 Posamentier, H.W., Ross, C.A. & Kendall, C.G.St.C (eds) *Sea-level Changes: An*  
1093 *Integrated Approach*. SEPM, Special Publication, **42**, 125-154.  
1094  
1095 Posamentier, H.W. & Kolla, V. 2003. Seismic Geomorphology and Stratigraphy of  
1096 Depositional Elements in Deep-Water Settings. *Journal of Sedimentary Research*, **73** (3),  
1097 367-388. DOI: 10.1306/111302730367  
1098  
1099 Provan, D.M.J. 1976. Composite Well Log 17/11-2. A/S Norske Shell.  
1100 [http://www.npd.no/engelsk/cwi/pbl/wellbore\\_documents/338\\_02\\_Composite\\_well\\_log.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/338_02_Composite_well_log.pdf)  
1101  
1102  
1103 Rattey, R.P. & Hayward, A.B. 1993. Sequence stratigraphy of a failed rift system: the  
1104 Middle Jurassic to Early Cretaceous basin evolution of the Central and Northern North  
1105 Sea. *In*: Parker, J.R. (ed) *Petroleum Geology: North-West Europe and Global Perspectives*  
1106 *- Proceedings of the 4th Petroleum Geology Conference*. Geological Society, London,  
1107 Petroleum Geology Conference series, **4**, 215–249. DOI: 10.1144/0040215  
1108  
1109 Ray, F.M., Pinnock, S.J., Katamish H. & Trunbull, J.B. 2010. The Buzzard Field: anatomy  
1110 of the reservoir from appraisal to production. *In*: Vining, B.A. & Pickering, S.C. (eds)  
1111 *Petroleum Geology: From Mature Basins to New Frontiers – Proceedings of the 7th*  
1112 *Petroleum Geology Conference*. Geological Society, London, Petroleum Geology  
1113 Conference series, **7**, 369-386. DOI: 10.1144/0070369  
1114  
1115 Rossland, A., Escalona, A. & Rolfsen, R. 2013. Permian-Holocene tectonostratigraphic  
1116 evolution of the Mandal High, Central Graben, North Sea. *AAPG Bulletin*, **97**, 923–957.  
1117 DOI: 10.1306/12031212093  
1118  
1119 Sears, R.A., Harbury, A.R., Protoy, A.J.G. & Stewart, D.J. 1993. Structural styles from the  
1120 Central Graben in the UK and Norway. *In*: Parker, J.R. (ed) *Petroleum Geology: North-*  
1121 *West Europe and Global Perspectives - Proceedings of the 4th Petroleum Geology*  
1122 *Conference*. Geological Society, London, Petroleum Geology Conference series, **4**, 1231-  
1123 1243. <https://doi.org/10.1144/0041231>  
1124  
1125 Skibeli, M., Barnes, K., Straume, T., Syvertsen, S.E. & Shanmugam, G. 1995. A sequence  
1126 stratigraphic study of Lower Cretaceous deposits in the northernmost North Sea. *In*: Steel,  
1127 R.J. (ed) *Sequence Stratigraphy on the Northwest European Margin*. NPF Special  
1128 Publication, **5**, 389-400.  
1129  
1130 Statoil. 1991. Well completion report Amalie-1 and Amalie 1A. Statoil Efterforskning OG  
1131 Produktion A/S, Copenhagen, Denmark.  
1132  
1133 Stewart, S.A. 2007. Salt tectonics in the North Sea Basin: a structural style template for  
1134 seismic interpreters. *In*: Ries, A.C., Butler, R.W.H., & Graham, R.H. (eds) *Deformation of*

- 1135 *the Continental Crust: The Legacy of Mike Coward*. Geological Society, London, Special  
1136 Publications, **272**, 361–396. DOI: 10.1144/GSL.SP.2007.272.01.19
- 1137
- 1138 Strass, I.F. 1979. Lithology. Well 7/3-1. *NPD Paper*, **20**.  
1139 [http://www.npd.no/engelsk/cwi/pbl/NPD\\_papers/164\\_01\\_NPD\\_Paper\\_No.20\\_Lithology\\_](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/164_01_NPD_Paper_No.20_Lithology_)  
1140 [\\_Well\\_7\\_3\\_1.pdf](http://www.npd.no/engelsk/cwi/pbl/NPD_papers/164_01_NPD_Paper_No.20_Lithology_)
- 1141
- 1142 Surlyk, F., Dons, T., Clausen, C.K. & Higham, J. 2003. Upper Cretaceous. *In*: Evans, D.,  
1143 Graham, C., Armour, A. & Bathurst, P. (eds) *The Millennium Atlas: Petroleum Geology of*  
1144 *the Central and Northern North Sea*. Geological Society, London, 213–233.
- 1145
- 1146 Thorsrud, A., Jensen, J.R. & Rasmussen, B.L. 2002. Final Well Report – Well Svane-  
1147 1&1A.
- 1148
- 1149 UKDD (UK Discovery Digest). 2007. Triassic Reservoirs of the Central Graben. Canadian  
1150 Discovery Ltd., **9**, 181–189. <http://ukdigest.canadiandiscovery.com/?q=node/35892>
- 1151
- 1152 Underhill, J.R. & Partington, M.A. 1993. Jurassic thermal doming and deflation in the  
1153 North Sea: implications of the sequence stratigraphic evidence. *In*: Parker, J.R. (ed)  
1154 *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 4th*  
1155 *Petroleum Geology Conference*. Geological Society, London, Petroleum Geology  
1156 Conference series, **4**, 337–345. DOI: 10.1144/0040337
- 1157
- 1158 Van Wijhe, D.H., 1987. Structural evolution of inverted basins in the Dutch offshore.  
1159 *Tectonophysics*, **137**, 171–219. Doi: [https://doi.org/10.1016/0040-1951\(87\)90320-9](https://doi.org/10.1016/0040-1951(87)90320-9)
- 1160
- 1161 Van Winden, M., de Jager, J., Jaarsma, B. & Bourollec, R. This Volume. New insights in  
1162 salt tectonics in the northern Dutch offshore: a framework for exploration. *In*: Kilhams, B.,  
1163 Kukla, P., Mazur, S., McKie, T., Mijnlief, H. & Van Ojik, K. (eds) Mesozoic resource  
1164 potential of the Southern Permian Basin. Geological Society of London Special  
1165 Publications.
- 1166
- 1167 Vejrbæk, O.V, Andersen, C., Duser, M., Herngreen, G.F.W., Krabbe, H., Leszczyński,  
1168 Lott, G.K., Mutterlose, J. & Van Der Molen, A.S. 2010. Cretaceous. *In*: Doornenbal, J.C.  
1169 and Stevenson, A.G. (eds) *Petroleum Geological Atlas of the Southern Permian Basin*  
1170 *Area*. EAGE Publications b.v., Houten, 194–209.
- 1171
- 1172 Veeken, P. & Rauch-Davies, M. 2006. AVO attribute analysis and seismic reservoir  
1173 characterisation. *First Break*, **2**, 7–18.  
1174 DOI: 10.3997/1365-2397.2006004
- 1175
- 1176 Verolles, F. 1982. Geological completion report. Well 3/7-3. Elf Aquitaine Norge A/S  
1177 Exploration Division. 311D/82/79-R.  
1178 [http://www.npd.no/engelsk/cwi/pbl/wellbore\\_documents/293\\_03\\_3\\_7\\_3\\_Completion\\_Rep](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/293_03_3_7_3_Completion_Report.pdf)  
1179 [ort.pdf](http://www.npd.no/engelsk/cwi/pbl/wellbore_documents/293_03_3_7_3_Completion_Report.pdf)

- Verweij, J.M. & Witmans, N. 2009. Terschelling Basin and southern Dutch Central Graben. Mapping and Modelling – Area 2A. *TNO Report*, **TNO-034-UT-2009-01569**.
- Voigt, S., Wagreich, M., Surlyk, F., Walaszczyk, I., Ulinčný, D., Čech, S., Voigt, T., Wiese, F., Wilmsen, M., Niebuhr, B., Reich, M., Funk, H., Michalík, Jagt, J.W.M., Felder, P.J. & Schulp, A. 2008. Cretaceous. In: McCann, T. (ed) *Geology of Central Europe*. Geological Society, London, 923–998.  
DOI: 10.1144/CEV2P.3
- Vollset, J. & Doré, A.G. 1984. A revised Triassic and Jurassic nomenclature for the Norwegian North Sea. *NPD Bulletin*, **3**.  
<http://www.npd.no/Global/Norsk/3-Publikasjoner/NPD-Bulletin/Bulletinenr3.pdf>
- Walker, J.D., Geissman, J.W., Bowring, S.A. & Babcock, L.E. 2012. GSA Geologic Time Scale v. 4.0. Geological Society of America.  
<https://www.geosociety.org/documents/gsa/timescale/timescl.pdf>
- Wong, Th.E. 2007. Jurassic. In: Wong, Th.E., Batjes, D.A.J. & De Jager, J. (eds) *Geology of the Netherlands*. Royal Netherlands Academy of Science and Art, Amsterdam, 241–264.
- Wong, Th.E., Van Doorn, H.M. & Schroot, B.M. 1989. »Late Jurassic« petroleum geology of the Dutch Central North Sea Graben. *Geologische Rundschau*, **78** (1), 319–336.  
doi:10.1007/BF01988367
- Zanella, E., Coward, M.P., & McGrandle, A. 2003. Structural Framework. In: Evans, D., Graham, C., Armour, A. & Bathurst, P. (eds) *The Millennium Atlas: Petroleum Geology of the Central and Northern North Sea*. Geological Society, London, 44–60.

1215 **TABLE CAPTION**


1216 Table 1. List of sources for well data

Type of data	Well	Datasource
Lithostratigraphic tops for seismic interpretation	Norway	NPD 2017, Shell database
	UK	Shell database
	Denmark	Shell database
Well shown in well panel Fig. 6	UK 30/11b-1, UK 29/5a-5	Milton-Worssell <i>et al.</i> 2006
	NO 2/1-8	Fjellanger 1986; NPD 2013
	NO 2/7-15	Phillips 1981; Isaksen & Tonstad 1989
	NO 3/7-3	Verolles 1982; Boirie & Jeannou 1984; NPD 2017
	Amalie-1	Statoil 1991
Other well data described in text and other images	Norway	NPD 2017
	NO 7/3-1	NPD 1979a; Strass 1979
	NO 17/10-1	NPD 1979b; Olsen 1979
	NO 17/11-1	A/S Norske Shell 1968
	NO 17/11-2	Provan 1976
	UKCG	Milton-Worssell <i>et al.</i> 2006
	Denmark (general)	Shell database
	Sten-1	Kern <i>et al.</i> 1983
	Tabita-1	Bonde <i>et al.</i> 1994
	Iris-1	Britoil 1985; Bonde <i>et al.</i> 1994
	Svane	Thorsrud <i>et al.</i> 2002
	West Lulu-4	Mærsk 1987
BCU and Top Lower Cretaceous subcrop data	e.g. NO 7/11-8, NO 3/7-1	NPD 2017; Shell database

1217

1218


1219 **FIGURE CAPTIONS**



1220


1221 **Fig. 1. (a)** Structural map of the Late Jurassic Central Graben depicting the study area  
1222 (Norwegian Central Graben area and parts of the UK Central Graben) and the adjacent UK  
1223 Central Graben study area of Milton-Worsell *et al.* (2006), that in combination define the  
1224 extended study area. **(b)** Detailed map of the study area, indicating seismic coverage (blue)  
1225 and the extent of available seismic depth maps (thick dotted outline). Dotted lines indicate  
1226 interpreted seismic sections S1-S6 (Fig. 5). CBH: Cleaver Bank High, DCG: Dutch Central  
1227 Graben, EOH: East Orkney High, NDB: Norwegian-Danish Basin, ECG: East Central  
1228 Graben, FG: Feda Graben, FGS: Fladen Ground Spur, FMH: Forties-Montrose High, HH:  
1229 Halibut High, IMF: Inner Moray Firth, JH: Jæren High, MNSH: Mid North Sea High, MT:  
1230 Marnock Terrace, OMF: Outer Moray Firth, RFH: Ringkøbing-Fyn High, SB: Søgne Basin,  
1231 SH: Sørvestlandet High, SG: Step Graben, SGH: Schill Ground High, TEG: Tail End Graben,  
1232 UH: Utsira High, WCG: West Central Graben, WGG: With Ground Graben, WP: Western  
1233 Platform. Modified after Fraser *et al.* (2003), Milton-Worsell *et al.* (2006) and Pharaoh *et al.*  
1234 (2010).






**Fig. 2.** Norwegian lithostratigraphy for the study area (left) and overview of Lower Cretaceous reservoirs in the Central North Sea (right). CrKn: Cromer Knoll Group, FB: Fischerbank Schiefer, NS: North Sea, SPB: Southern Permian Basin. Seismic horizon abbreviations from top to bottom: BC: Base Cenozoic, TLC: Top Lower Cretaceous, BCU: Base Cretaceous Unconformity, TMJ: Top Middle Jurassic, TZ: Top Zechstein, TR: Top Rotliegend. Modified after Vollset & Doré (1984), Van Wijhe (1987), Isaksen & Tonstad (1989), Wong *et al.* (1989), Copestake *et al.* (2003), Milton-Worsell *et al.* (2006), De Jager & Geluk (2007), Jakobsen *et al.* (2005); Herngreen & Wong (2007), UKDD (2007), Wong (2007), NDD (2012). Geological timescale dates after Walker *et al.* (2012).






**Fig. 3.** Time depth maps of four regional horizons in the study area: (a) Base Cenozoic; (b) Top Lower Cretaceous; (c) Base Cretaceous Unconformity (BCU); (d) Top Rotliegend. Note that the Base Cenozoic time depth map (a) is also the Cenozoic isochron map and that the Top Rotliegend map is incomplete due to locally poor seismic quality. Dotted lines indicate the trace of interpreted transects S1-S6 and white dots indicate well locations along these transects (see Fig. 5). Solid outlines indicate the extent of the 3D seismic survey. Dashed outlines indicate the extent of the available previously interpreted seismic horizons in the UK and Denmark (see Fig. 1). BB: Breiflabb Basin, CT: Cod Terrace, FG: Feda Graben, HH: Hydra High, JoH: Josephine High; FMH: Forties-Montrose High; MNSH: Mid North Sea High, LR: Lindsess Ridge, PT: Pigvarr Terrace, RFH: Ringkøbing-Fyn High, SH: Sørvestlandet High, ST: Steinbit Terrace.




**Fig. 4.** Isochron maps showing the location and geometry of depocenters in the study area in Two-way travel time. **(a)** UC: Upper Cretaceous isochron map; **(b)** LC: Lower Cretaceous (Cromer Knoll Group). Dotted lines indicate the trace of interpreted transects S1-S6 and wells along these transects (see Fig. 5). Solid outlines indicate the extent of the 3D seismic survey. Dashed outlines indicate the extent of the available previously interpreted seismic horizons in the UK and Denmark (see Fig. 1). The larger yellow dots indicate >3 m sand occurrences in wells within the study area, whereas crosses indicate sandstone traces (<3 m thickness). Wells: (1) UK 29/5a-5, (2) UK 30.11b-1, (3) NO 2/1-8, (4) NO 2/7-15, (5) NO 3/7-3, (6) Sten-1, (7) Amalie-1, (8) Tabita-1, (9) Iris-1, (10) Svane-1. BB: Breiflabb Basin, CT: Cod Terrace, FG: Feda Graben, FMH: Forties-Montrose High, HH: Hydra High, JoH: Josephine High, MH: Mandal High, MNSH: Mid North Sea High, LR: Lindessness Ridge, PT: Pigvarr Terrace, RFH: Ringkøbing-Fyn High, SB: Søgne Basin, SH: Sørvestlandet High.

*Zwaan, Lower Cretaceous reservoirs, North Sea Central Graben*



**Fig 5.** Interpreted seismic sections S1-S3. UK: United Kingdom. For section locations see Fig. 1b. Reference datum is mean sea level.







**Fig 5. (continued)** Interpreted seismic sections S4-S6. UK: United Kingdom. For section locations see Fig. 1b. Section S6 modified after Møller & Rasmussen (2003). Reference datum is mean sea level.



1285  
1286  
1287  
1288  
1289  
1290




**Fig. 7.** Interpretation of reservoir potential in the extended study area for the Latest Ryazanian (BCU-level). Well data, possible sand source areas, Ryazanian fault activity, possible sediment transport fairways and areas of possible sandstone development are projected on top of the Lower Cretaceous isochron map. Well identifiers are Ir: Iris-1, St: Sten-1, Sv: Svane-1, Ta: Tabita-1, (1): well NO 3/7-1. AR: Auk Ridge, ArF: Argyll Field, CT: Cod Terrace, FBB: Fisher Bank Basin, FFT: Flora-Fife Trend, FMH: Forties-Montrose High, GG: Gertrud Graben, HH: Hydra High, IH: Inge High, JH: Jæren High, JoH: Josephine High, MH: Mandal High, MT: Marnock Terrace, PT: Piggvar Terrace, RFH: Ringkøbing-Fyn High, SB: Søgne Basin, SBB: South Buchan Basin, SH: Sørvestlandet High, TEG: Tail End Graben, WP: Western Platform. Modified after Japsen *et al.* (2003), Milton-Worssell *et al.* (2006) and Rossland *et al.* (2013).




**Fig. 8.** Interpretation of reservoir potential throughout the extended study area for the Aptian-Albian (near-Top Lower Cretaceous level). Well data, possible sand source areas, interpreted possible sediment transport fairways and areas of possible sandstone development are projected on top of the Lower Cretaceous isochron map. AR: Auk Ridge, ArF: Argyl Field CT: Cod Terrace, FBB: Fisher Bank Basin, FF: Flora Field, FMH: Forties-Montrose High, GG: Gertrud Graben; HH: Hydra High, IH: Inge High, JH: Jæren High, JoH: Josephine High, MH: Mandal High, MT: Marnock Terrace, PT: Piggvar Terrace; RFH: Ringkøbing-Fyn High, SB: Søgne Basin, SBB: South Buchan Basin, SH: Sørvestlandet High, TEG: Tail End Graben, WP: Western Platform. Modified after Japsen *et al.* (2003), Milton-Worsell *et al.* (2006) and Rossland *et al.* (2013).






**Fig. 9.** Proposed Early Cretaceous paleogeographic situation around the Mandal High area. Image modified after Rossland *et al.* (2013).





**Fig. 10.** Idealised cross section proposing the main potential reservoirs and traps in the Mandal High-Søgne Basin area. C: Cenozoic, UC: Upper Cretaceous, LC: Lower Cretaceous, UJ: Upper Jurassic. LJ: Lower Jurassic, Tr: Triassic, Z: Zechstein (evaporites), R: Rotliegend, B: Pre-Permian sediments and/or (metamorphic) Basement. Image modified after Rossland *et al.* (2013).



1329

1330

**Fig. 11. (a)** Gross depositional environment overview of the Central and Southern North Sea in Ryazanian times (K10). **(b)** Gross depositional environment of the South Viking Graben area in Aptian times (K40-50), corresponding to the Aptian-Albian reservoir interval. Lower Cretaceous sand presence: (1) Edvard Grieg field, (2) Johan Sverdrup field, (3) well NO 7/10-1, (4) well NO 7/11/-1, (5) well NO 7/11-2, (6) well NO 7/3-1, ÅG: Åsta Graben, BF: Britannia Field, BFB: Broad Fourteens Basin, CBH: Cleaver Bank High, COS: Central Offshore Saddle, CS: Crawford Spur, CVG: Central Viking Graben, DCG: Dutch Central Graben, EOH: East Orkney High, ESP: East Shetland Platform, FP: Friesland Platform, HH: Halibut High, IMF: Inner Moray Firth, JH: Jæren High, FMH: Forties-Montrose High, KF: Kopervik Fairway, MH: Mandal High, MT: Marnock Terrace, NCG: Norwegian Central Graben, NDB: Norwegian-Danish Basin, OMF: Outer Moray Firth, RFH: Ringkøbing-Fyn High, RVG: Roer Valley Graben, SG: Step Graben, SGH: Schill Grund High, SPB: Sole Pit Basin, TB: Terschelling Basin, TEG: Tail End Graben (Danish Central Graben), UH: Utsira High, UKCG: UK Central Graben, WGG: Witch Ground Graben, WNB: West Netherlands Basin. Modified after Copestake *et al.* (2003), NPD (2017) for the South Viking Graben area, after Milton-Worsell *et al.* (2006), Copestake *et al.* (2003), Rossland *et al.* (2013) for the UKCG and NCG, after Vejbæk *et al.* (2010), after Pharaoh *et al.* (2010) for the Danish, German and (parts of) the Dutch Central Graben, and after Jeremiah *et al.* (2010) for the Southern North Sea.